

Diagnosis and Management of Cancer Risk in the Gastrointestinal Hamartomatous Polyposis Syndromes: Recommendations From the US Multi-Society Task Force on Colorectal Cancer

C. Richard Boland, MD^{1,*}, Gregory E. Idos, MD, MS^{2,*}, Carol Durno, MD^{3,4}, Francis M. Giardiello, MD⁵, Joseph C. Anderson, MD, MHCDS, FACG^{6,7,8}, Carol A. Burke, MD, FACG⁹, Jason A. Dominitz, MD, MHS^{10,11}, Seth Gross, MD, FACG¹², Samir Gupta, MD^{13,14,15}, Brian C. Jacobson, MD, MPH, FACG¹⁶, Swati G. Patel, MD, MS¹⁷, Aasma Shaukat, MD, MPH, FACG^{18,19}, Sapna Syngal, MD, FACG^{20,21,22} and Douglas J. Robertson, MD, MPH^{6,7}

The gastrointestinal hamartomatous polyposis syndromes are rare, autosomal dominant disorders associated with an increased risk of benign and malignant intestinal and extraintestinal tumors. They include Peutz-Jeghers syndrome, juvenile polyposis syndrome, the *PTEN* hamartoma tumor syndrome (including Cowden's syndrome and Bannayan-Riley-Ruvalcaba syndrome), and hereditary mixed polyposis syndrome. Diagnoses are based on clinical criteria and, in some cases, confirmed by demonstrating the presence of a germline pathogenic variant. The best understood hamartomatous polyposis syndrome is Peutz-Jeghers syndrome, caused by germline pathogenic variants in the *STK11* gene. The management is focused on prevention of bleeding and mechanical obstruction of the small bowel by polyps and surveillance of organs at increased risk for cancer. Juvenile polyposis syndrome is caused by a germline pathogenic variant in either the *SMAD4* or *BMPRI1A* genes, with differing clinical courses. Patients with *SMAD4* pathogenic variants may have massive gastric polyposis, which can result in gastrointestinal bleeding and/or protein-losing gastropathy. Patients with *SMAD4* mutations usually have the simultaneous occurrence of hereditary hemorrhagic telangiectasia (juvenile polyposis syndrome–hereditary hemorrhagic telangiectasia overlap syndrome) that can result in epistaxis, gastrointestinal bleeding from mucocutaneous telangiectasias, and arteriovenous malformations. Germline pathogenic variants in the *PTEN* gene cause overlapping clinical phenotypes (known as the *PTEN* hamartoma tumor syndromes), including Cowden's syndrome and related disorders that are associated with an increased risk of gastrointestinal and colonic polyposis, colon cancer, and other extraintestinal manifestations and cancers. Due to the relative rarity of the hamartomatous polyposis syndromes, recommendations for management are based on few studies. This US Multi-Society Task Force on Colorectal Cancer consensus statement summarizes the clinical features, assesses the current literature, and provides guidance for diagnosis, assessment, and management of patients with the hamartomatous polyposis syndromes, with a focus on endoscopic management.

Am J Gastroenterol 2022;117:846–864. <https://doi.org/10.14309/ajg.000000000001755>

The gastrointestinal hamartomatous polyposis syndromes are rare, autosomal dominant disorders associated with an increased risk of benign and malignant intestinal and extraintestinal tumors. Nevertheless, there has been tremendous progress in recent years, both in understanding the underlying genetics that underpin these disorders and in elucidating the biology of associated premalignant and malignant conditions. Emerging data in

affected populations focus increasingly on those with defined cancer susceptibility germline pathogenic variants leading to less heterogeneity in terms of quantifying cancer risk.

The US Multi-Society Task Force on Colorectal Cancer (USMSTF) is a group of colorectal cancer (CRC) content experts appointed by the American College of Gastroenterology, American Gastroenterological Association, and American Society for

¹Division of Gastroenterology, University of California-San Diego School of Medicine, San Diego, California, USA; ²Divisions of Gastroenterology and Clinical Cancer Genomics, Center for Precision Medicine, City of Hope National Medical Center, Duarte, California, USA; ³The Zane Cohen Centre for Digestive Diseases, Mount Sinai Hospital, Toronto, Ontario, Canada; ⁴Division of Gastroenterology, Hepatology and Nutrition, Hospital for Sick Children, University of Toronto, Toronto, Ontario, Canada; ⁵Division of Gastroenterology, Department of Medicine, The Johns Hopkins University School of Medicine, Baltimore, Maryland, USA; ⁶Veterans Affairs Medical Center, White River Junction, Vermont, USA; ⁷Geisel School of Medicine at Dartmouth, Hanover, New Hampshire, USA; ⁸University of Connecticut, Farmington, Connecticut, USA; ⁹Department of Gastroenterology, Hepatology and Nutrition, Cleveland Clinic, Cleveland, Ohio, USA; ¹⁰Veterans Affairs Puget Sound Health Care System, Seattle, Washington, USA; ¹¹University of Washington School of Medicine, Seattle, Washington, USA; ¹²Division of Gastroenterology and Hepatology, New York University Langone Health, New York, New York, USA; ¹³Veterans Affairs Medical Center, San Diego, California, USA; ¹⁴University of California San Diego, La Jolla, California, USA; ¹⁵Veterans Affairs San Diego Healthcare System, San Diego, California, USA; ¹⁶Division of Gastroenterology, Massachusetts General Hospital, Boston, Massachusetts, USA; ¹⁷University of Colorado School of Medicine, Rocky Mountain Regional Veterans Affairs Medical Center, Aurora, Colorado, USA; ¹⁸Minneapolis Veterans Affairs Health Care System, Minneapolis, Minnesota, USA; ¹⁹University of Minnesota, Minneapolis, Minnesota, USA; ²⁰Brigham and Women's Hospital, Boston, Massachusetts, USA; ²¹Dana-Farber Cancer Institute, Boston, Massachusetts, USA; ²²Harvard Medical School, Boston, Massachusetts, USA. **Correspondence:** C. Richard Boland, MD, E-mail: crboland@health.ucsd.edu.

This article is being published jointly in *The American Journal of Gastroenterology*, and *Gastrointestinal Endoscopy*. The article is identical except for minor stylistic and spelling differences in keeping with each journal's style. Citations from any of the 3 journals can be used when citing this article.

*Authors share co-first authorship.

Received March 23, 2022; accepted March 24, 2022; published online April 26, 2022

Gastrointestinal Endoscopy, supplemented at times by other experts to complement existing expertise. In this USMSTF Consensus Statement, the gastrointestinal hamartomatous polyposis syndromes were chosen because of recent progress in understanding these diseases. The following entities reviewed are: Peutz-Jeghers syndrome (PJS), juvenile polyposis syndrome (JPS), *PTEN* hamartoma tumor syndrome (PHTS, including Cowden's syndrome [CS] and Bannayan-Riley-Ruvalcaba syndrome [BRRS]), and hereditary mixed polyposis syndrome (HMPS). Germline alterations are known to cause each of these disorders, but the diagnosis can also be made on the basis of clinical criteria.

Although there are essentially no long-term prospective controlled studies of comparative effectiveness of management strategies for these syndromes, there have been consensus statements by expert panels that made management recommendations for these disorders. The goal of this USMSTF Statement was to review the literature focusing on the most recent data, synthesize both the data and the suggested approaches to diagnosis and management by other expert groups, and present the consensus recommendations of the USMSTF (Table 1). Review of summary tables, conference calls, and revisions of iterative drafts, including recommendation statements, were used to reach consensus, at which time documents were forwarded to Governing Boards for approval. As our USMSTF is a group of individuals with expertise in gastroenterology and gastrointestinal malignancies, we have reserved our management recommendations to these areas and defer to other expert groups' recommendations for other cancers (which are reviewed here). This document therefore recommends clinical approaches to diagnosing and managing these conditions that affect children and adults, focuses on cancer risk, and provides insights into future research opportunities.

METHODS

A computer-aided PubMed search was performed from 2000 to 2018, with additional back searches as required, and consisted of the following search terms: *hamartoma*, *hamartomatous polyp*, *hamartoma syndrome*, *Peutz-Jeghers syndrome*, *juvenile polyp*, *juvenile polyposis*, *Cowden's syndrome*, *Cowden's disease*, *PTEN-hamartoma*, *Bannayan-Riley-Ruvalcaba syndrome*, *hyperplastic polyposis*, *serrated polyposis*, and *hereditary mixed polyposis syndrome*. Only English-language articles were reviewed. Published articles were selected on the basis of relevance to the diagnosis or clinical management of these diseases. Emphasis was placed on the risk for gastrointestinal cancer in these disorders to gain consensus on rational and reasonable strategies for management once these diseases are diagnosed in a family. The document was approved by the governing boards of each of the sponsoring gastroenterology societies.

The USMSTF approach to an adapted use of Grading of Recommendations Assessment, Development and Evaluation (GRADE) has been described previously.¹ In brief, the GRADE process categorizes the quality of the evidence as high, moderate, low, or very low on the basis of the strength of underlying studies, and that categorization can be adjusted on the basis of study limitations. For example, randomized trials begin as high-quality evidence and observational studies as low-quality evidence, but their quality may be adjusted up or down on the basis of specific study factors. Although

Table 1. Questions and Recommendations of Best Practice

Which individuals with hamartomatous polyps should be referred for genetic evaluation?

We recommend patients with any of the following undergo a genetic evaluation: 2 or more lifetime hamartomatous polyps, a family history of hamartomatous polyps, or a cancer associated with a hamartomatous polyposis syndrome in first or second-degree relatives. Genetic testing (if indicated) should be performed using a multigene panel test. (Strong recommendation, low quality of evidence)

Peutz-Jeghers syndrome

Who should undergo a genetic evaluation for Peutz-Jeghers syndrome?

We recommend genetic evaluation for any individual with the following: 1) 2 or more histologically confirmed Peutz-Jeghers polyps, 2) any number of Peutz-Jeghers polyps in an individual who has a family history of Peutz-Jeghers syndrome in a first-degree relative, 3) characteristic mucocutaneous pigmentation in a person with a family history of Peutz-Jeghers syndrome, 4) any number of Peutz-Jeghers polyps in a person with the characteristic mucocutaneous pigmentation of Peutz-Jeghers syndrome. (Strong recommendation, low quality of evidence)

Which organs should undergo surveillance when caring for a patient with Peutz-Jeghers syndrome?

Patients with Peutz-Jeghers syndrome are at increased risk for cancer in multiple organs including cancer of the breast, small bowel, colon, stomach, pancreas, ovaries, testes, and lungs.

Given this risk, we recommend a multidisciplinary approach to cancer surveillance in these organs (Strong recommendation, low quality of evidence)

How and when should small bowel surveillance be performed in Peutz-Jeghers syndrome?

We recommend that baseline small bowel surveillance using video capsule endoscopy or magnetic resonance enterography be performed between ages 8-10 years or earlier if the patient is symptomatic. If no polyps are found at the initial examination, surveillance should resume at age 18. Because of the risk of small bowel intussusception, small bowel surveillance in adulthood is recommended to continue throughout life every 2-3 years. (Strong recommendation, low quality of evidence)

What is the recommended approach to endoscopic surveillance of the colon, stomach, and duodenum in Peutz-Jeghers syndrome?

We suggest a baseline upper gastrointestinal endoscopy between the ages of 8 and 10 years, which could be performed at the time of capsule placement for small bowel surveillance or if polyps are identified on magnetic resonance enterography. Although the initiation age for colonoscopy remains uncertain, we also suggest initiation of colonoscopy at same time as esophagogastroduodenoscopy. In those in whom characteristic polyps are detected, both colonoscopy and esophagogastroduodenoscopy should be repeated every 2-3 years. In those in whom there are no Peutz-Jeghers polyps at baseline, surveillance is repeated at age 18 years, or sooner should symptoms arise, and then every 3 years. (Weak recommendation, very low quality of evidence)

What size polyps found on small bowel imaging in Peutz-Jeghers syndrome should be removed?

We recommend polypectomy of small bowel polyps that are symptomatic or ≥ 10 mm to prevent intussusception and other complications, such as bleeding. (Strong recommendation, low quality of evidence)

What is the recommended pancreatic cancer surveillance in Peutz-Jeghers syndrome?

We suggest annual pancreatic cancer surveillance with either magnetic resonance cholangiopancreatography or endoscopic ultrasound starting at age 35 years. (Weak recommendation, low quality of evidence)

Table 1. (continued)**Juvenile Polyposis Syndrome**

Who should undergo a genetic evaluation for juvenile polyposis syndrome? We recommend genetic evaluation for any individual with 1) 5 or more juvenile polyps of the colon or rectum; or 2) 2 or more juvenile polyps in other parts of the gastrointestinal tract; or (3) any number of juvenile polyps and 1 or more first-degree relatives with juvenile polyposis syndrome. (Strong recommendation, low quality of evidence)

Which organs should undergo surveillance when caring for a patient with juvenile polyposis syndrome?

Juvenile polyposis syndrome patients are at increased risk for cancer in multiple organs including cancer of the colon and stomach.

Given this risk, we recommend patients with juvenile polyposis syndrome undergo surveillance of the colon and stomach. (Strong recommendation, low quality of evidence)

At what age should colonoscopic and upper endoscopic surveillance begin in individuals identified with juvenile polyposis syndrome?

We suggest initiating colonoscopic and upper endoscopic surveillance at age 12–15 years, or earlier if symptomatic. Surveillance should be repeated every 1–3 years depending on polyp burden. (Weak recommendation, low quality of evidence)

Which patients with juvenile polyposis syndrome should undergo screening for hereditary hemorrhagic telangiectasia?

We suggest patients with SMAD4 pathogenic variants be clinically evaluated for HHT at the time of the diagnosis, including screening for and appropriate management of cerebral and pulmonary AVMs. (Weak recommendation, low quality of evidence)

PTEN hamartoma tumor syndrome

Which gastrointestinal findings should prompt a genetic evaluation for *PTEN* hamartoma tumor syndrome?

We recommend individuals with multiple gastrointestinal hamartomas or ganglioneuromas undergo genetic evaluation for Cowden's syndrome and related conditions. (Strong recommendation, low quality of evidence)

Which organs should undergo surveillance for cancer when caring for a patient with *PTEN* hamartoma tumor syndrome?

In *PTEN* hamartoma tumor syndrome, patients are at increased risk for cancer in multiple organs, including cancer of the breast, thyroid, kidney, uterus, colon, and skin.

Given this risk, we recommend a multi-disciplinary approach to cancer surveillance in these organs. (Strong recommendation, low quality of evidence)

What is the recommended colonoscopic surveillance in individuals identified with *PTEN* hamartoma tumor syndrome?

We suggest colonoscopy surveillance to begin at age 35 years (or 10 years younger than age of any relative with colorectal cancer), repeated at intervals no greater than 5 years, depending on polyp burden. (Weak recommendation, low quality of evidence)

NOTE. Specific circumstances may merit modification of the recommendations. In cases where very-early-onset cancers develop, the above statements may be modified to start surveillance 10 years earlier than the youngest cancer diagnosis in the family.

the GRADE process entails a formal meta-analysis to assess the quality of evidence for each recommendation, the USMSTF employs a modified, qualitative approach for this assessment. The GRADE process separates evaluation of the quality of the evidence to support

a recommendation from the strength of that recommendation. This is done in recognition of the fact that, although the quality of the evidence can influence the strength of the recommendation, other factors can influence a recommendation, such as adverse effects, patient preferences, values, and cost. Generally, strong recommendations mean that most informed patients would choose the recommended management. Weak recommendations mean that patients' choices will vary according to their values and preferences, and clinicians should ensure that patient care is in keeping with their values and preferences. When the quality of the evidence to support a recommendation is low or very low, or if there is a close balance between desirable and undesirable consequences, then usually only a weak recommendation would be warranted. Weaker recommendations are indicated by phrases such as "we suggest," and stronger recommendations are typically stated as "we recommend."

However, the relative infrequency of, and absence of controlled prospective trials of the interventions (eg, to prevent cancer) in, these syndromes leave all of the recommendations without a robust basis of underlying evidence. Thus, all of the interventional recommendations fall (at best) into the "low quality of evidence" GRADE category, indicating that the true effect of the interventions may be markedly different than estimated at this time, and that further research is likely to impact or change our confidence in the effects. As such, this review is intended to establish a starting point for future research into the care of patients with the hamartomatous polyposis syndromes.

Cancer Family History Assessment and Referral for Genetic Testing in Gastroenterology Practice

Question: Which individuals with hamartomatous polyps should be referred for genetic evaluation?

Recommendation: We recommend patients with any of the following undergo a genetic evaluation: 2 or more lifetime hamartomatous polyps, a family history of hamartomatous polyps, or a cancer associated with a hamartomatous polyposis syndrome in first- or second-degree relatives. Genetic testing (if indicated) should be performed using a multigene panel test. (Strong recommendation, low quality of evidence)

Hereditary cancer syndromes account for approximately 5%–10% of new cancer diagnoses and many of the cancers that arise in families with undiagnosed hereditary cancer syndromes are preventable. The identification of individuals with a hereditary gastrointestinal cancer syndrome requires a thorough evaluation of the patient's personal and family history of cancer. The collection and assessment of family cancer history is a valuable tool for cancer interception and prevention and can be critical in the identification of genetic susceptibility. An accurate family history is one that collects the following information: 1) type of cancer, 2) age at diagnosis of each primary cancer, 3) lineage (maternal or paternal), 4) ethnicity (people of some ethnicities, such as those with Ashkenazi Jewish ancestry, are at greater risk for certain cancers), and 5) results of any previous cancer-related genetic testing.²

Features of a patient's personal history and clinical characteristics may suggest an inherited susceptibility to cancer. Although it is not rare to identify individuals with an isolated hamartomatous polyp (particularly an isolated juvenile polyp), other features may prompt further evaluation for an underlying hereditary syndrome. Features associated with




Figure 1. Histologic characteristics of polyps. hamartomatous polyps. Images courtesy of Drs Aaron Pollett and Thomas Plesec. (A) Peutz-Jeghers polyp: pathognomonic broad bands of mucosal smooth muscle seen throughout the lesion. (B) Juvenile polyp: characteristic chronic inflammatory infiltrate and cystic dilatation. (C) Adenomatous polyp: characteristic hypercellularity with glandular crowding, enlarged nuclei, increased mitotic activity and reduced goblet cells. By definition, all tubular adenomas show epithelial dysplasia. (D) Ganglioneuroma: benign neuroectodermal tumor composed of ganglion and Schwann cells.

the hamartomatous polyposis syndromes are outlined in detail in this document and include early age at cancer onset, multiple cancers in close relatives, unusual numbers of hamartomatous polyps, or associated dermatologic findings.

Genetic evaluation may include genetic counseling and/or genetic testing.

Genetic counseling is a key component to hereditary cancer risk assessment. The purpose of genetic counseling is to educate




Figure 2. Endoscopic images of hamartomatous polyposis syndromes. Endoscopic photos provided courtesy of Swati Patel, MD, Gregory Idos, MD, and Carol Burke, MD. (A) Peutz-Jeghers small bowel polyps. (B) Peutz-Jeghers gastric polyps. (C) Peutz-Jeghers colon polyps. (D) Juvenile polyposis gastric polyps. (E) Juvenile polyposis colon polyp. (F) Cowden syndrome associated—esophageal glycogenic acanthosis. (G) Cowden syndrome small bowel polyps. (H) Cowden syndrome colon polyps. (I) Cowden syndrome gastric polyps.

individuals about the genetic and biologic factors that are related to a patient’s cancer diagnosis or risk of disease. Counseling helps an individual understand the relevant genetic, medical, and psychosocial information to make informed

decisions about their health care. This includes reviewing and expanding the following: family history information, elements of genetic testing, tailored cancer risks associated with a pathogenic variant, impact on medical management, reproductive

issues and options, confidentiality of results, risks with genetic discrimination, potential significance of test results for other family members, and other pertinent topics. Practice guidelines from the American College of Medical Genetics and Genomics and National Society of Genetic Counselors are available for details regarding the elements and process of genetic counseling.³ Although traditional models of genetic evaluation and testing included a certified genetic counselor, alternative models exist and are emerging that include provisions for pretest counseling to be provided by physicians and other health care providers in order to deal with the increasing demand for genetic testing. If a patient is found to be a carrier of a germline pathogenic variant, or the results are ambiguous due to the finding of a variant of uncertain significance, the help of a genetics provider for post-test counseling and education is recommended. In the current era, the vast majority of genetic testing for inherited cancer risk predisposition is performed using a multigene panel testing approach.⁴ Some patients or families may elect to decline genetic testing due to concerns about risk to confidentiality and insurance; in these cases, surveillance may still be indicated in the presence of a concerning clinical and/or family history.

If a germline pathogenic variant is identified, other family members should be offered testing for clarification of their own risk. This testing may facilitate initiation of screening for associated cancers before symptomatic manifestations occur and reduce the morbidity and mortality associated with the syndrome. For example, early small bowel surveillance may find a polyp that could be removed from a child with a pathogenic variant in *STK11* before leading to intussusception. It is important to recognize that genetic testing may not identify pathogenic variants in every family suspected of a hereditary syndrome. However, there may be clinical features in the family history that suggest a familial predisposition to cancer and suggest more intensive surveillance recommendations. Referral to Centers of Excellence might be particularly helpful when genetic testing results are ambiguous in the setting of suspicious features and prophylactic surgery is being considered. Lastly, in the era of multigene panel testing, there may be a scenario in which a germline variant is found incidentally associated with an unsuspected syndrome. In these cases, patients may be eligible for cancer screening and surveillance as outlined. However, phenotype and cancer risk compared with patients with classic familial features are not established and are areas of active research. Enlisting the assistance of a genetic specialist may be particularly helpful in interpreting ambiguous results and providing management recommendations in these cases.

When children are identified with a hamartomatous polyposis syndrome, their transition of care to adulthood for cancer surveillance is a unique aspect that bears consideration. It is imperative to transition adolescents with life-long medical conditions from child-centered to adult-centered care. Preparation for this transition takes place throughout childhood and adolescence to achieve independent health management in adulthood. Steps required are individualized based on the developmental needs of the patient. Inherited conditions involve generational factors, as multiple family members may be affected. Health care providers can assist in the transition of care by coordinating screening and surveillance to ensure patients receive recommended care.⁵

GASTROINTESTINAL HAMARTOMATOUS POLYPOSIS SYNDROMES

The hamartomatous polyposis syndromes are rare entities with an estimated prevalence of 1/100,000–200,000,^{6,7} but this has not been measured directly in any population. The term *hamartoma* implies a non-neoplastic tumor with a markedly distorted architecture composed of an abnormal mixture of cells and tissue normally present in that particular area. The diagnosis is based on the presence of a pathogenic germline variant or meeting clinical criteria for the syndrome. The hamartomatous polyposis syndromes are distinct from Lynch syndrome and the adenomatous polyposis syndromes, based on the presence of hamartomas (Figures 1 and 2). Certain hamartomatous polyps of the gut have a unique histopathological appearance, such as those associated with PJS, PHTS, JPS, and HMPS.⁸ Hamartomas are not typically characterized by dysplasia, but some evidence suggests the existence of a hamartoma–carcinoma pathway in some of these polyps.

PEUTZ-JEGHERS SYNDROME

Question: Who should undergo a genetic evaluation for Peutz-Jeghers syndrome?

Recommendation: We recommend genetic evaluation for any individual with the following: 1) 2 or more histologically confirmed Peutz-Jeghers polyps, 2) any number of Peutz-Jeghers polyps in an individual who has a family history of Peutz-Jeghers syndrome in a first-degree relative, 3) characteristic mucocutaneous pigmentation in a person with a family history of Peutz-Jeghers syndrome, and 4) any number of Peutz-Jeghers polyps in a person with the characteristic mucocutaneous pigmentation of Peutz-Jeghers syndrome. (Strong recommendation, low quality of evidence)

Clinical Features

PJS was the first hamartomatous polyposis syndrome described, by Peutz in Holland in 1921 and by Jeghers, McKusick, and Katz in the United States in 1949.⁹ The clinical recognition of PJS was facilitated by the characteristic mucocutaneous freckling around the mouth and multiple cerebriform-appearing polyps due to smooth muscle bands coursing through the polyp (Figures 1A and 2A–C, and Table 2). Hamartomatous polyps vary in size and may have a characteristic histologic structure, which makes it possible to distinguish the PJ polyp. PJ polyps are typically composed of branching bands of smooth muscle covered by hyperplastic glandular mucosa.¹⁰ PJS polyps may develop in the stomach, small intestine, and colon. Rectal bleeding with anemia is the most common presentation, followed by abdominal pain, diarrhea, and intussusception. The clinical management in early life is initially focused on preventing complications of small bowel polyposis-related obstruction and bleeding and, in adulthood, the focus is primarily on management of cancer risk.

Diagnosis

The diagnostic clinical features of PJS include the presence of 2 or more histologically confirmed PJ polyps; any number of PJ polyps in an individual who has a family history of PJS in a first-degree relative; characteristic mucocutaneous pigmentation in a person with a family history of PJS; or any number of PJ polyps in a person with the characteristic mucocutaneous pigmentation of PJS.¹¹

Table 2. Clinical Features of Gastrointestinal Hamartomatous Polyposis Syndromes

Syndrome	Commercially available gene testing	Polyps	Clinical features
PJS	<i>STK11</i>	Peutz-Jeghers polyps (pathologically characteristic)	Childhood: Labial pigmentation; gastrointestinal bleeding and intussusception Adults: Increased risk for multiple cancers
JPS	<i>SMAD4</i> or <i>BMPR1A</i>	Juvenile (inflammatory) polyps; juvenile polyps and inflammatory polyps are pathologically indistinct	Childhood: Gastrointestinal bleeding, auto-amputation of polyps; anemia Adults: CRC and gastric cancer
CS	<i>PTEN</i> (inactivation) <i>WWP1</i> (gain-of-function)	Hyperplastic polyps; juvenile-like polyps; ganglioneuromas; lipomas; hamartomas; adenomas	Childhood: none Adults: multiple cancer risks
BRRS	<i>PTEN</i>	Same as for CS	Developmental delay, hemangiomas, lipomas, gastrointestinal polyps
HMPS	<i>GREM1</i> (duplication upstream of promoter)	Pathologically mixed with features of adenoma, hyperplastic polyps, inflammatory polyps	Increased risk of colonic polyposis and CRC

Genetics

In 1997, a genetic locus for PJS was mapped to chromosome 19p13.3,¹² which in 1998 led to the cloning of the *STK11* (serine/threonine kinase) gene, which encodes the LKB1 (liver kinase B1) protein, and linkage to PJS.¹³ *STK11* functions like a tumor suppressor gene, regulates cell growth via adenosine monophosphate-activated protein kinase,¹⁴ and negatively regulates mTOR signaling.¹⁵

PJS is inherited in an autosomal dominant fashion with an inactivating germline pathogenic variant inherited from the affected parent. It was initially assumed that the polyps occurred after the loss of the second, wild-type, allele inherited from the unaffected parent in a somatic tissue according to the classic “two-hit” model of Knudson.¹⁶ However, loss of the wild-type allele (ie, the second hit) is not an obligatory feature of PJ polyps.¹⁷ Moreover, recent data in mice indicate that the presence of a single inactivating germline pathogenic variant (ie, haploinsufficiency), as occurs in individuals with PJS, promotes the development of gastrointestinal polyposis, and that loss of the wild-type allele is not necessary for the formation of the polyps in these mice. Furthermore, conditional knockout of the *STK11* gene targeted to gastrointestinal smooth muscle cells yields the same polyposis phenotype in mice.¹⁸ Evidence suggests that LKB1 deficiency in either T cells¹⁹ or mesenchymal cells^{18,19} leads to immune cell proliferation in the stroma. Stromal deficiency of LKB1 leads to tumor formation (in mice) via the interleukin-11–JAK/STAT3 (Janus kinase/signal transducer and activator of transcription 3) pathway, and administration of the JAK1/2 inhibitor ruxolitinib dramatically reduces polyposis in mice.²⁰ The phenomenon of stromal-driven epithelial polyp development may be referred to as a “landscaper” genetic effect.²¹ These discoveries raise the possibility of the development of novel preventive pharmacological interventions for patients.

Genotype–phenotype relationships reveal that missense alterations in *STK11* are associated with later onset of symptoms than other sequence variations.^{11,22} Some of the germline pathogenic variants in *STK11* are *Alu*-mediated deletions and inversions.²³ Deletions and inversions are likely to completely inactivate a gene, whereas a missense pathogenic variant may have an intermediate effect on protein function. Approximately 15% of *STK11* pathogenic variants in PJS involve large genetic deletions.²⁴ Therefore, genetic diagnostic platforms must include strategies to detect these types of germline

variations; at least some of the “missing” germline pathogenic variants may involve obscure rearrangements of the *STK11* gene.

Cancer Risk

Very high lifetime risks of cancer occur in multiple organs in patients with PJS, inside and outside the gut. Intratubular large-cell hyalinizing Sertoli cell neoplasms, ovarian sex cord tumors, and adenoma malignum of the uterine cervix, although not common, are linked to PJS²⁵ (Table 3).

The lifetime risk of colon, stomach, and small bowel cancer has been estimated at 12%–39%, 29%, and 13% respectively^{6,26–28} (Table 3). Most CRCs occur after the mid-20s (range, 27–71 years, median age 46 years), but these malignancies have been reported in teenage years as well.²⁶ Mean age at diagnosis of gastric adenocarcinoma ranges between 30 and 40 years and of small intestinal cancer between 37 and 42 years.^{26,27,29}

In a meta-analysis of 210 cases reported in 6 publications (retrospective cohort studies with kindred-based ascertainment from the United States, United Kingdom, and The Netherlands) on PJS that were based on clinical and histologic criteria with varying types of ascertainment, the relative risk (RR) for any cancer was 15.2 (95% confidence limits [CL], 2 and 19) compared with a variety of time period–specific US-based registries. Significantly increased age-adjusted cancer risks were noted for the small intestine (RR, 520; 95% CL, 220 and 1306; cumulative risk [CR], 13%), stomach (RR, 213; 95% CL, 96 and 368; CR, 29%), pancreas (RR, 132; 95% CL, 44 and 261; CR, 36%), colon (RR, 84; 95% CL, 47 and 137; CR, 39%), esophagus (RR, 57; 95% CL, 2.5 and 557; CR, 0.5%), ovary (RR, 27; 95% CL, 7.3 and 68; CR, 21%), lung (RR, 17.0;

Question: Which organs should undergo surveillance when caring for a patient with Peutz-Jeghers syndrome?

Recommendation: Patients with Peutz-Jeghers syndrome are at increased risk for cancer in multiple organs, including cancer of the breast, small bowel, colon, stomach, pancreas, ovaries, testes, and lungs. Given this risk, we recommend a multidisciplinary approach to cancer surveillance in these organs. (Strong recommendation, low quality of evidence)

Table 3. Risk of Cancer in Hamartomatous Polyposis Syndromes

Site	General population risk, ^a %	Syndrome risk, %	Mean age at diagnosis, y	Reference
PJS				
Colorectal	4.3	39	42–46	26, 29
Stomach	<1	29	30–40	26, 29
Small bowel	<1	13	37–42	26, 29
Breast	12.9	32–54	37–59	26, 28, 29
Ovarian (mostly SCTAT)	1.2	21	28	26
Cervix (adenoma malignum)	<1	10–23	34–40	26
Uterus	3.1	9	43	26, 29
Pancreas	1.7	11–36	41–52	26, 28, 29, 32
Testicular (Sertoli cell tumor)	<1	9	6–9	26, 29
Lung	6.3	7–17	47	26, 28, 29
JPS				
Colon	4.3	39	44	86
Stomach	<1	5–21	54	65, 67, 89
CS				
Breast	12.9	25–85	38–46	100, 101, 102
Thyroid	1.3	3–38	31–38	100, 101, 102
Uterus	3.1	5–28	25	95, 100, 101, 102
Kidney (renal cell)	1.7	15–34	40	97, 100, 104
Colon	4.3	9–18	35	100–103, 106
Melanoma	2.3	6	3 ^b	100, 101
HMPS				
Colon	4.3	Increased	—	119–121
SCTAT, sex cord tumors with annular tubules.				
^a National Cancer Institute. Surveillance, Epidemiology, and End Results Cancer Statistics Review 1975–2017. Lifetime risk (%) of being diagnosed with cancer by site, 2017.				
^b Youngest age of onset.				

95% CL, 5.4 and 39; CR, 15%), uterus (RR, 16.0; 95% CL, 1.9 and 56; CR, 9%), testes (RR, 4.5; 95% CL, 0.12 and 25; CR, 9%), and breast (RR, 15.2; 95% CL, 7.6 and 27; CR, 54%).²⁶ Mean age at the time of cancer diagnosis was 42.9 years. The absolute risk of developing any cancer between the ages of 15 and 64 was estimated to be 93%.

The risk of cancer in PJS was revisited in 2 follow-up studies that included some who were members of the original cohort.^{28,30} The lattermost report included some of the original 210 cases and expanded to 419 cases, in which 297 had confirmed germline pathogenic variants in *STK11*.²⁸ The cumulative incidences of cancer by decade from age 20 to 70 years were 2%, 5%, 17%, 31%,

60%, and 85% respectively, confirming the initial estimates and tumor spectrum—predominantly the gastrointestinal tract and breast. Cancer risks were the same in those with a clinical diagnosis but no detectable germline pathogenic variant in the *STK11* gene.

Several collaborative studies from Europe and the United States again found similar risks for cancer.^{29,31} A systematic review of 20 cohort studies looking at 1644 patients with PJS confirmed at least 1 cancer in >90% of patients with PJS at a mean age of 42 years.²⁹ CRC was the most commonly diagnosed tumor, followed by cancers of the breast, small intestine, stomach, lung, pancreas, cervix, ovary, bile ducts, and testicles. A multicenter study from Italy of 119 *STK11* pathogenic variant carriers reported an overall RR for cancer of 22.0 in women (in part because of additional risks for cervical cancers) and 8.6 in men, compared with an Italian-based general population registry, and a cumulative risk of cancer reaching 89% by age 65 years.²⁷ A more recent study from China confirmed elevated cancer risks (albeit somewhat lower than the American and European data), early age of onset, and a similar tumor spectrum.³²

Surveillance of Affected Individuals

The effectiveness of cancer surveillance in PJS has not been evaluated in controlled studies. Consequently, surveillance recommendations have been developed by consensus groups and expert opinion analyzing cancer risks and published organ-specific surveillance protocols (Tables 3 and 4).^{11,33–35}

Gastrointestinal Polyposis and Cancer

Gastric, small bowel, and colorectal polyposis occur in 88%–100% of patients, with the majority appearing in the small bowel (60%–90%) and colon (50%–64%).³⁶ Polyps can vary in number (1–100) and size (0.1–3 cm in diameter) and age of onset of symptoms may vary. Polyp growth begins in childhood by age 10 years (33%), with most experiencing symptoms such as bleeding, abdominal pain, intussusception, or obstruction (68%) by age 18 years.³⁷ In affected or at-risk individuals, early surveillance with esophagogastroduodenoscopy (EGD), colonoscopy, and small bowel imaging with video capsule endoscopy (VCE) and/or magnetic resonance enterography (MRE) is recommended to begin at age 8 years.^{11,33–35}

Intussusception is rare in children younger than 5 years, and the precise risk of intussusception between 5 and 18 years of age is unknown. Retrospective registry data report that 23 of 34 adults with PJS (68%) had undergone laparotomy by the age of 18 years, 70% of which were performed as an emergency. By the age of 10 years, 30% had required a laparotomy.³⁷ In a single-institution

Question: How and when should small bowel surveillance be performed in Peutz-Jeghers syndrome?

Recommendation: We recommend that baseline small bowel surveillance using video capsule endoscopy or magnetic resonance enterography be performed between ages 8 and 10 years or earlier if the patient is symptomatic. If no polyps are found at the initial examination, surveillance should resume at age 18 years. Because of the risk of small bowel intussusception, small bowel surveillance in adulthood is recommended to continue throughout life every 2–3 years. (Strong recommendation, low quality of evidence)

Table 4. Surveillance Guidelines in Hamartomatous Polyposis Syndromes

Examination	ACG 2015		NCCN 2020		ESPGHAN 2019		USMSTF 2020		Evidence grade
	Age of initiation, y	Interval, y	Age of initiation, y	Interval, y	Age of initiation, y	Interval, y	Age of initiation, y	Interval, y	
JPS									
Colonoscopy	12–15	1–3	15	2–3	12–15	—	12–15	1–3	Low
Upper endoscopy	12–15	1–3	15	2–3	Late teens ^a	—	12–15	1–3	Low
Screen for vascular lesions	≤6 mo	—	≤6 mo	—	At diagnosis	—	≤6 mo ^b	—	—
PJS									
Colonoscopy	8, 18 ^c	3	Late teens	2–3	8	3	8–10, 18 ^c	2–3	Very low
Upper endoscopy	8, 18 ^c	3	Late teens	2–3	8	3	8–10, 18 ^c	2–3	Very low
VCE	8, 18 ^c	3	~8–10 ^d	2–3	8	3	8, 18 ^c	2–3	Low
CT or MRE of small bowel	—	—	~8–10 ^d	2–3	—	—	—	—	—
MRI/MRCP or EUS of pancreas	30	1–2	~30–35 ^e	1–2	—	—	35	1	Low
MRI and/or mammogram	25	1	~25	1	—	—	—	—	—
Physical examination	Birth to teenage	1	~10	1	—	—	—	—	—
Pelvic examination and Pap smear	25 ^f	1	~18–20	1	—	—	—	—	—
Testicular examination	Birth to teenage	1	~10	1	—	—	—	—	—
PHTS									
Colonoscopy	15	2	35	5	—	—	35	5	Low
Upper endoscopy	15	2–3	—	—	—	—	—	—	—
Thyroid examination and US	Adolescence	1	7	1	—	—	—	—	—
MRI and/or mammogram	30–35	1	30–35	1	—	—	—	—	—
Endometrial sampling	30–35	1	—	1–2	—	—	—	—	—
Urinalysis or renal US	18	1	40	1–2	—	—	—	—	—
Skin examination	~18	1	At diagnosis	1	—	—	—	—	—

ACG, American College of Gastroenterology; CT, computed tomography; ESPGHAN, European Society for Paediatric Gastroenterology Hepatology and Nutrition; EUS, endoscopic ultrasound; MRCP, magnetic resonance cholangiopancreatography; NCCN, National Comprehensive Cancer Network.

^aUpper gastrointestinal screening in JPS *SMAD4* carriers is indicated in asymptomatic patients starting in late teens. For non-*SMAD4* JPS patients, upper endoscopic screening is only indicated if the patient has relevant symptoms or anemia not explained by colonic polyps.

^b*SMAD4* mutation carriers should be clinically evaluated for HHT at the time of the diagnosis, including screening for, and appropriate management of, cerebral and pulmonary AVMs.

^cFirst procedure at 8 years of age; if polyps present, repeat every 3 years; if no polyps, restart at 18 years of age and every 3 years.

^dBaseline at age 8–10 years of age with follow-up interval based on findings but at least by age 18 years of age, then every 2–3 years, although this may be individualized by at least age 18 years, then every 2–3 years, although this may be individualized, or with symptoms).

^eBased on clinical judgment, early initiation age may be considered, such as 10 years younger than the earliest age of onset in the family.

^fACG 2015 Guidelines recommend transvaginal ultrasound as part of surveillance beginning at age 25 years.

study of 379 pediatric patients who underwent pneumatic reduction for intussusception (from all causes), one-quarter required operative management.³⁸ There is a paucity of studies that assess modalities to evaluate the small bowel in patients with PJS. Retrospective data comparing VCE to small bowel barium studies have reported VCE as a useful diagnostic tool in PJS. A retrospective study from France included 27 children who underwent at least 1 VCE.³⁹ The authors found VCE was a useful diagnostic tool, however, findings at VCE may not predict future bowel obstructions. Although VCE may not predict future bowel

obstructions, it has a greater sensitivity in detecting small bowel polyps compared with MRE and provides patients with another alternative if they are unable to undergo MRE. Ten children with PJS with polyps >15 mm identified by VCE and MRE underwent single-balloon enteroscopy.⁴⁰ In this small study, single-balloon enteroscopy was effective for treating small bowel polyps. Further larger, multicenter studies are warranted to accurately determine the safety of therapeutic single-balloon enteroscopy in children.⁷ Based on the published data, VCE or MRE is recommended between ages 8 and 10 years in asymptomatic patients.

Investigations should commence earlier if patients have symptoms. If no polyps are found on the baseline investigation, repeat small bowel evaluation may commence at age 18 years. If polyps are found, further investigation and surveillance should be individualized based on polyp size and location. Because of the risk of small bowel intussusception, small bowel surveillance in adulthood continues throughout life every 2–3 years. If polyps are present in the colon or stomach, EGD or colonoscopy is continued as necessary every 2–3 years.

Management of Polyposis

Although the malignant potential of PJ polyps is unknown and the evidence of benefit from gastrointestinal surveillance is not robust, endoscopic removal of polyps is recommended to prevent polyp-related complications and to reduce the risk of cancer. In a study of long-term outcomes of gastrointestinal procedures from a PJ polyposis registry, investigators tracked the results of 776 procedures among 63 patients with PJS at a median age of 20 years (range, 3–59 years). A total of 2461 polypectomies were performed; more than 1000 polyps were removed during colonoscopy and more than 400 polyps were removed during EGD, and the remaining polyps were removed by means of enteroscopy or laparotomy of the small bowel.⁴¹ A substantial proportion of patients required intervention for removal of large polyps and the authors concluded that surveillance reduced polyp burden and likelihood of polyp-related complications, and provided cancer surveillance. Therefore, we recommend polypectomy for polyps in the stomach and colorectum that are >0.5 cm in size on endoscopic surveillance and an attempt to remove all polyps if endoscopically feasible.^{33,35,42,43}

In the small bowel, balloon enteroscopy and MRE have similar diagnostic yields for lesions ≥ 15 mm, but endoscopy permits polyp removal.⁴³ Consequently, removal, preferably by enteroscopy, of small intestinal polyps that are symptomatic or rapidly growing, or asymptomatic polyps that are >1.0–1.5 cm in size, has been recommended.^{33,35,37,42,44,45} Surgery is often needed when small bowel intussusception occurs. Some authorities recommend an attempt to clear the small intestine of polyps during laparotomy by means of intraoperative endoscopy with polypectomy or, for larger polyps, by means of enterotomy. This “clean sweep” approach appears to decrease the need for recurrent small bowel surgery.⁴⁶ Thus, the early management focus is on large hamartomatous polyps and their tendency to obstruct the gut or bleed. As children grow older and transition into adulthood, the focus shifts to managing the cancer risk.

Question: What is the recommended approach to endoscopic surveillance of the stomach, duodenum, and colon in Peutz-Jeghers syndrome?

Recommendation: We suggest a baseline upper gastrointestinal endoscopy between the ages of 8 and 10 years, which could be performed at the time of capsule placement for small bowel surveillance or if polyps are identified on magnetic resonance enterography. Although the initiation age for colonoscopy remains uncertain, we also suggest initiation of colonoscopy at the same time as esophagogastroduodenoscopy. In those in whom characteristic polyps are detected, both colonoscopy and esophagogastroduodenoscopy should be repeated every 2–3 years. In those in whom there are no Peutz-Jeghers polyps at baseline, surveillance is repeated at age 18 years, or sooner should symptoms arise, and then every 3 years. (Weak recommendation, very low quality of evidence)

Breast Cancer

Invasive ductal adenocarcinoma poses the greatest risk of early malignancy to patients with PJS (24%–54% lifetime risk in women) and often presents at a young age (range, 19–61 years in 1 study).^{26,47} The risk of breast cancer in women with PJS is within the same range as women affected by *BRCA-1* or *BRCA-2* pathogenic variants (40%–85% lifetime risk for breast cancer).^{26,48} Therefore, consensus opinion surveillance recommendations by groups that include breast cancer experts^{49,50} include monthly breast self-examination starting at age 18 years, biannual clinical breast examination starting at age 25 years, annual breast magnetic resonance imaging (MRI) from ages 25–29 years, and mammography with consideration of tomosynthesis (3-dimensional mammography) alternating every 6 months with breast MRI with contrast from ages 30 to 75 years.^{6,11} The option of prophylactic mastectomy might be discussed on a case-by-case basis, factoring in family history. In patients with PJS, referral to a breast cancer specialist for management of breast cancer surveillance is reasonable, and a multidisciplinary approach including a breast surgeon is recommended when prophylactic mastectomy is being considered.

Question: What size polyps found on small bowel imaging in Peutz-Jeghers syndrome should be removed?

Recommendation: We recommend polypectomy of small bowel polyps that are symptomatic or ≥ 10 mm to prevent intussusception and other complications, such as bleeding. (Strong recommendation, low quality of evidence)

Pancreas Cancer

Pancreatic cancer is the third most frequently occurring malignancy in patients with PJS. The lifetime risk of pancreatic ductal adenocarcinoma is between 11% and 36%, presenting, on average, at ages 41–52 years, with 95% of cases occurring after age 24 years.^{6,26,28,51,52} PJS is associated with the highest relative risk of all pancreatic cancer syndromes, with the exception of hereditary pancreatitis (25%–40% lifetime risk).⁴³ Consequently, a high-risk pancreatic surveillance protocol is recommended. The consensus recommendations of the International Cancer of the Pancreas Screening Consortium⁵³ recommend pancreas MRI/magnetic resonance cholangiopancreatography and/or endoscopic ultrasound every 1–2 years starting from age 40 years, for which evidence suggests that these cancers can be found in earlier stages.⁵⁴ The National Comprehensive Cancer Network Guidelines recommend initiating pancreatic cancer surveillance between the ages of 30 and 35 years. Due to reports of pancreatic cancer diagnoses in patients with PJS before the age of 40 years,²⁶ the USMSTF suggests initiating annual surveillance at age 35 years with MRI/magnetic resonance cholangiopancreatography and/or endoscopic ultrasound. Ideally, these examinations would alternate on an annual basis, as they are complementary. In addition, routine fasting glucose and hemoglobin A1c at initiation of screening is recommended by the Cancer of the Pancreas Screening Consortium. At the current time, the USMSTF awaits definitive data before making a recommendation regarding fasting glucose and hemoglobin A1c. Recent updates to the Cancer of the Pancreas Screening Guidelines detail surveillance protocols and management recommendations based on imaging and endoscopic findings.⁵⁵

Question: What is the recommended pancreatic cancer surveillance in Peutz-Jeghers syndrome?

Recommendation: We suggest annual pancreatic cancer surveillance with either magnetic resonance cholangiopancreatography or endoscopic ultrasound starting at age 35 years. (Weak recommendation, low quality of evidence)

Gynecological Cancers

The lifetime risks for ovarian, uterine, and cervical cancer are estimated at 21%, 9%, and 10%–23%, respectively, with mean age at presentation of 28–35 years, 43 years, and 34–40 years, respectively.^{6,26,27} Of note, almost all ovarian tumors in patients with PJS are sex cord tumors with annular tubules, and rarely cystadenomas or granulosa cell tumors. Sex cord tumors with annular tubule neoplasms are a heterogeneous group of benign or malignant neoplasms, but the tumors rarely have lymph node metastasis.²⁶ Also, an unusual percentage of cervical cancers in patients with PJS are adenoma malignum, a rare, well-differentiated, cervical adenocarcinoma that is associated with a poor prognosis and difficult to diagnose on Pap smear. A high level of suspicion is required for diagnosis.⁵⁶ Recommendations for gynecological surveillance are pelvic examination with Pap smear and transvaginal ultrasound annually starting at age 25 years.^{33,35}

Testicles

The estimated risk of testicular cancer in male patients with PJS is 9%, mean age at diagnosis is 9 years (range, 3–20 years).^{6,26,27} These tumors present as testicular masses. The tumors are Sertoli cell tumors that can cause gynecomastia and other signs of hyperestrogenism and occasionally virilization and/or accelerated height growth.²⁶ Accelerated height velocity can be challenging to detect as adolescents have growth spurts as part of normal maturation and development. Expert opinion recommends annual history and physical examination (including self-examination) with observation for feminizing changes and examination of the testicles^{6,34,35}; based on the range of age at diagnosis of this tumor, examination should start from birth. Ultrasound of the testicles every 2 years from birth to age 12 years has been suggested.⁴⁴

Lung

The lifetime risk of lung cancer in patients with PJS has been estimated between 7% and 17%,^{6,26–28,30} compared with 0.2%–0.6% in nonsmokers in the general population.⁵⁷ The cumulative risk of lung cancer in PJS surpasses 5% by age 55 years.⁶ Of note, lung cancer risk in patients with PJS has not been calculated with adjustment for smoking status. The RRs of lung cancer in patients with PJS compared with nonsmokers are similar to people with a more than 30 pack-year history of smoking who have quit for 10–15 years (hazard ratio, 14.8).⁵⁸ Currently, the American College of Chest Physicians, American Society of Clinical Oncology, and American Cancer Society recommend low-dose computed tomography annually for individuals with this level of cumulative risk for lung cancer from ages 55 to 74 years.^{59,60} There are no data to show benefit of lung cancer surveillance in patients with PJS. Lung cancer surveillance with annual low-dose computed tomography, as performed in smokers at high risk for lung cancer, may be considered in patients with PJS. Smoking cessation counseling in patients with PJS is advisable to mitigate risk.^{33,34}

Chemoprevention

Currently, there are no known chemopreventive agents in clinical practice that slow or prevent the development of intestinal polyps and cancers in PJS. Pathogenic variants in the *STK11* gene decrease inhibition of mTOR leading to the development of intestinal polyps. A trial examining the oral selective mTOR inhibitor, everolimus, was stopped prematurely because of poor patient accrual. Only 2 patients were enrolled, 1 with pancreatic cancer that progressed and another patient who withdrew from the protocol because of severe complications from the medication.⁶¹

In a murine model of PJS, treatment with celecoxib, a COX2 inhibitor, resulted in a >50% reduction in polyp burden. When used in patients with PJS with diffuse polyposis in the body of the stomach (tens to hundreds), 2 of 6 had a significant reduction in polyp number, as assessed by 5 independent evaluators, after administering celecoxib (200 mg twice per day for 6 months).⁶²

Summary

PJS is associated with a very high cumulative lifetime risk of cancer of multiple organs, including, but not limited to, the gastrointestinal tract. Intensive surveillance is recommended to prevent and manage complications of polyposis and identify cancer at an early stage. The development of polyposis and the cancer risks may be a reflection of the effects of haploinsufficiency of the LKB1 protein, for which there are possible medical therapies to be explored. It is unclear whether the basic mechanisms responsible for hamartoma formation in younger life are the same as those that create the risks for cancer later in life. Simulation models and clinical trials are needed to optimize endoscopic surveillance frequencies and the use of imaging modalities in adults. In addition, collaborative multicenter consortia may help facilitate chemoprevention trials in the future.

JUVENILE POLYPOSIS SYNDROME

Question: Who should undergo a genetic evaluation for juvenile polyposis syndrome?

Recommendation: We recommend genetic evaluation for any individual with 1) 5 or more juvenile polyps of the colon or rectum; 2) 2 or more juvenile polyps in other parts of the gastrointestinal tract; or 3) any number of juvenile polyps and 1 or more first-degree relatives with juvenile polyposis syndrome. (Strong recommendation, low quality of evidence)

Clinical Features

JPS is an autosomal dominant inherited condition in which multiple juvenile polyps are found in the colorectum (98% of affected patients) (Figure 1B), stomach (14%), jejunum and ileum (7%), and duodenum (7%).^{63–65} The incidence of JPS is between 1 in 100,000 and 1 in 160,000 individuals.⁶⁵ The polyps in JPS vary in size from small sessile nodules to pedunculated lesions that are ≥ 3 cm in diameter. Histologically, the typical juvenile polyp has a distinctive cystic architecture, dilated mucus-filled glands, a prominent lamina propria, and Paneth cells enmeshed within a dense infiltration of inflammatory cells. In patients with germline pathogenic variants in *SMAD4*, additional features of gastric polyposis, gastric cancer, and a JPS–hereditary hemorrhagic telangiectasia (HHT) overlap syndrome are common. The

management of JPS is based on decreasing the risk of bleeding and gastric and colorectal cancer through polypectomy. Patients with JPS-HHT overlap syndrome should have lifelong HHT surveillance. One study also demonstrated that as many as 38% may have thoracic aorta abnormalities.⁶⁶

Juvenile polyps are most often solitary and are not syndromic, occurring sporadically in infants and children. A typical history associated with a solitary juvenile polyp is the asymptomatic passing of a polyp into an infant's diaper. Juvenile polyps appear endoscopically to have a smooth red surface, may be sessile or pedunculated, and pathologically are characterized by cystic dilatation of mucus-filled glands suspended in an inflamed stroma (Figures 1B and 2D and E). These lesions may also be called retention polyps or inflammatory polyps because of their microscopic appearance.

Diagnosis

The diagnosis of JPS is made based on clinical criteria or identification of a germline pathogenic variant in *SMAD4* or *BMPRIA*. The clinical diagnosis of JPS is made when a person has any 1 of the following: 1) 5 or more juvenile polyps of the colon or rectum; 2) any number of juvenile polyps in parts of the gastrointestinal tract other than the colon; or 3) any number of juvenile polyps and 1 or more first-degree relatives with JPS.⁶⁵

Genetics

JPS is an autosomal dominant disorder with approximately 75% of cases inherited from a parent and 25% representing de novo pathogenic variants.⁶⁴ Approximately 60% of patients with JPS have a pathogenic variant in the *BMPRIA* or *SMAD4* gene.⁶⁷ Rarely (1 in 1,000,000) individuals develop features of JPS and PHTS, known as juvenile polyposis of infancy, due to a large deletion encompassing both the *BMPRIA* and *PTEN* genes.

In a study from the Cleveland Clinic of 35 patients with JPS, germline pathogenic variants in *SMAD4* and *BMPRIA* were associated with similar colonic polyposis phenotypes, but *SMAD4* pathogenic variant carriers were more likely to have more gastric polyps, and an 11% risk of gastrointestinal cancer.⁶⁸ However, no gastric cancers were reported in patients with *BMPRIA* pathogenic variants in 8 patients followed for a mean of 11 years.

However, germline pathogenic variants in other genes may cause a hamartomatous polyp phenotype. In a study of 49 patients referred to the Cleveland Clinic for unexplained hamartomatous or hyperplastic polyps, germline pathogenic variants were found in multiple genes, including *endoglin* (*ENG*, a gene associated with HHT), *STK11* (the PJS gene), *SMAD4*, *BMPRIA*, and *PTEN*.⁶⁹ In a later study from this group of 603 patients with a "moderate load of gastrointestinal polyps" with at least 1 confirmed to be a hamartoma or hyperplastic polyp, 13% were found to have a germline pathogenic variant in at least 1 of the genes listed above and 20% of the cohort had a personal history of CRC.⁷⁰

As discussed in the context of PJS, it has been traditionally thought that a germline pathogenic variant associated with a hereditary colon cancer syndrome did not change the biology of the normal tissues and a second (somatic) hit was required for a tumor to form. In fact, allelic loss of the *SMAD4* locus was found in the epithelial component of juvenile polyps of patients with JPS (together with a germline pathogenic variant in *SMAD4*) using *in situ* hybridization, but not in the inflammatory or stromal cells. This was compatible with the Knudson 2-hit model.⁷¹ However, another group has reported that *SMAD4* haploinsufficiency (ie, a

lower dose of the gene product due to the specific type of germline pathogenic variant) causes the JPS phenotype in humans, supported by data from mice.⁷² However, just as in PJS, being haploinsufficient for *SMAD4* (ie, just the germline pathogenic variant), is associated with partially diminished transforming growth factor- β signaling, at least in mice,⁷² as this alters proliferation in T cells, which contributes to the development of polyps and cancer.^{73,74} Specific deletion of *SMAD4* in mouse T cells leads to up-regulation of the Th17-inflammatory pathway in the stroma, and the growth of large polyps in the gastrointestinal tracts of mice.^{73,74} This raises the question of whether there is abnormal regulation of immune cells in JPS, and may explain the inflammatory appearance of juvenile polyps independent of abnormal biology in the epithelium—another landscaper mechanism, as discussed above for PJS. These observations may have important clinical implications for future attempts to halt the appearance or growth of the inflammatory polyps. However, allelic loss (ie, a somatic variants as the second hit) occurs in at least some polyps of JPS patients.^{75,76}

Juvenile Polyposis Syndrome–Hereditary Hemorrhagic Telangiectasia Overlap Syndrome

HHT occurs in approximately 15%–81% of patients with a germline *SMAD4* pathogenic variant.^{77,78} The clinical features of HHT, such as epistaxis, obscure gastrointestinal bleeding, digital clubbing, visceral arteriovenous malformations (AVMs), and mucocutaneous telangiectasias should be sought in *SMAD4* patients. International guidelines (outlined below) recommend that surveillance and treatment for HHT complications are necessary for all *SMAD4* carriers.⁷⁹

Juvenile Polyposis of Infancy

Juvenile polyposis of infancy is a severe form of juvenile polyposis. This disease presents in the first 2 years of life with diarrhea, abdominal pain, rectal bleeding, refractory anemia, hypoalbuminemia, and enteropathy. Case reports indicate that a large deletion in the long arm of chromosome 10 (10q23), encompassing the *PTEN* and *BMPRIA* genes, is associated with the development of the disease.^{69,80–83}

Gastrointestinal Polyposis and Cancer

In a retrospective chart review study of 257 children with juvenile polyps at a single-referral center, patients presented at a median age of 5.6 years, and at colonoscopy 39% had multiple polyps.⁸⁴ Among 192 patients who underwent complete colonoscopy at initial diagnosis, 117 (60.9%) had a single polyp and 75 (39.1%) had multiple polyps. These lesions recurred in 21 of 47 patients (44.7%) after an initial eradication, including 3 of 18 presenting with a single polyp, and neoplasia was found in 3.9% of lesions.

Question: Which organs should undergo surveillance when caring for a patient with juvenile polyposis syndrome?

Recommendation: Patients with juvenile polyposis syndrome are at increased risk for cancer in multiple organs, including cancer of the colon and stomach.

Given this risk, we recommend patients with juvenile polyposis syndrome undergo surveillance of the colon and stomach. (Strong recommendation, low quality of evidence)

Patients with JPS often have a variable presentation of polyp distribution, which may occur throughout the colon and/or stomach and the cancer risk is attributable to the presence of dysplasia in the polyps. One study of 78 juvenile polyps from 12 patients with JPS and 34 patients with sporadic juvenile polyps reported that dysplasia was present in 31% of the polyps from patients with JPS but in none of the polyps from patients with sporadic juvenile polyps.⁸⁵ Dysplasia in polyps from patients with JPS was associated with somatic variants in the *APC* gene.⁸⁵ In a longitudinal study from St Mark's Hospital of 44 patients with JPS from 30 kindreds, a total of 787 polyps (juvenile and adenomatous) were resected, and 65 of 787 (8.3%) contained mild/moderate architectural dysplasia, and 20 additional polyps (2.5%) were adenomatous.⁶⁵

Patients with JPS are at increased risk for cancer principally in the stomach and colon (Table 3). In a small cohort of patients with JPS ($n = 84$) relative to age-, sex-, and race-matched controls, the RR of CRC was estimated to be 34 (95% CL, 14.4 and 65.7), with a cumulative lifetime risk of CRC reaching 38.7%.⁸⁶ The CRCs were diagnosed at a mean age of 43.9 years. Interestingly, no other gastrointestinal cancers were noted in this cohort, but the numbers were small and the duration of the study was short. Individuals with gastric polyposis, usually in association with pathogenic variants in *SMAD4* are also at risk for gastric cancer, with the lifetime risk estimated to be at least 30% and median age of diagnosis is 58 years; this has not been reported in association with *BMPRIA* pathogenic variants.^{67,87-91} Estimates of gastrointestinal cancer risk range from 11%⁶⁸ to 55%,⁸⁹ but none of the studies are prospective long-term studies (which would lead to underestimates), and all are prone to referral-based ascertainment bias (overestimates).

Management of Polyposis

The goal of surveillance in JPS is to mitigate symptoms related to the disorder and decrease the risk of complications from the manifestations, including cancer. Colonoscopy should be first performed at age 12–15 years and repeated every 1–3 years, depending on polyp burden found, with removal of all polyps when feasible or at least all polyps ≥ 5 mm. Upper endoscopy should be first performed at age 12–15 years and repeated every 1–3 years depending on polyp burden found, with removal of polyps ≥ 5 mm. Due to the possible presence of intestinal telangiectasias, an annual history and physical examination and complete blood counts to monitor for rectal bleeding and/or anemia should begin at age 12–15 years in patients with a germline *SMAD4* pathogenic variant.

Endoscopic polypectomy is recommended for colorectal polyposis management. Surgery with colectomy and ileorectal anastomosis is recommended for patients with CRC, endoscopically unmanageable colon polyp burden and uncontrolled anemia from

colonic bleeding.³³ In some cases, proctocolectomy is necessary for rectal cancer or advanced polyp burden of the rectum. Colectomy in patients with JPS should be reserved for patients with polyp burdens that cannot be managed by polypectomy, persistent blood loss leading to severe anemia or hypoalbuminemia, or cancer. A decision to proceed to colectomy should be reviewed with gastroenterologists and surgeons with expertise in caring for individuals with hereditary polyposis syndromes.

The risk of gastric cancer is a concern in patients with JPS with *SMAD4* pathogenic variants and gastric polyposis. There is a paucity of published data evaluating the stomach and small bowel in pediatric patients with JPS. Those with upper gastrointestinal symptoms, or with anemia not explained by colonic polyps, should undergo evaluation with upper endoscopy.⁹² In pediatric patients without *SMAD4* pathogenic variants, according to the existing data, gastroscopy is not indicated unless the child has symptoms.⁹² In asymptomatic patients with *SMAD4* pathogenic variants, it is prudent to assess the upper tract between the ages of 12 and 15 years. At the time of this publication, it is uncertain as to whether *BMPRIA* pathogenic variants are associated with gastric cancer risk, and that pending new evidence, upper endoscopy surveillance is suggested at intervals similar to those recommended for *SMAD4* carriers. In adults, partial or complete gastrectomy is indicated in patients with gastric cancer, high-grade dysplasia, inability to adequately survey or endoscopically control polyposis, persistent anemia or gastrointestinal bleeding from gastric polyposis or angioectasia, symptoms of gastric outlet obstruction, or protein-losing gastropathy.^{63,64}

Management of Juvenile Polyposis Syndrome–Hereditary Hemorrhagic Telangiectasia

Question: Which patients with juvenile polyposis syndrome should undergo screening for hereditary hemorrhagic telangiectasia?

Recommendation: We suggest patients with *SMAD4* pathogenic variants be clinically evaluated for hereditary hemorrhagic telangiectasia at the time of the diagnosis, including screening for and appropriate management of cerebral and pulmonary arteriovenous malformations. (Weak recommendation, low quality of evidence)

As recommended by HHT Foundation International, patients with *SMAD4* pathogenic variants should be screened for vascular findings associated with HHT.^{79,93} Children with possible or confirmed HHT should be screened for brain AVMs at the time of diagnosis and undergo at least 1 follow-up MRI at puberty because brain AVM development appears to correlate with times of growth. Lung AVM screening and surveillance is recommended at diagnosis and then every 3–5 years with pulse oximetry testing and consideration of transthoracic contrast echocardiogram. In adulthood, surveillance should include annual hemoglobin or hematocrit for all patients older than 35 years. Transthoracic contrast echocardiogram as a screen for pulmonary AVMs should be performed at the time of diagnosis, within 5 years preceding planned pregnancy, after pregnancy, and otherwise every 5–10 years. Physicians should consider referring these patients to HHT Centers of Excellence for this evaluation. Brain MRI with and without contrast should be performed at birth or at the time of diagnosis to screen for cerebral vascular malformations.

Question: At what age should colonoscopic and upper endoscopic surveillance begin in individuals identified with juvenile polyposis syndrome?

Recommendation: We suggest initiating colonoscopic and upper endoscopic surveillance at age 12–15 years, or earlier if symptomatic. Surveillance should be repeated every 1–3 years depending on polyp burden. (Weak recommendation, low quality of evidence)

Chemoprevention

No known effective chemoprevention strategies exist for the development of polyposis in patients with JPS.

Summary

BMPRIA and *SMAD4* gene alterations are responsible for JPS, yet only 60% of patients will have a pathogenic alteration identified. The clinical overlap syndromes include JPS-HHT in patients with a *SMAD4* pathogenic variant and juvenile polyposis of infancy in patients with a combined *BMPRIA* and *PTEN* deletion. The symptoms of JPS are usually related to bleeding from colorectal polyposis or in *SMAD4*-related JPS to gastric polyposis or manifestations of HHT. Cancer risk in JPS is elevated, mainly in the colon and stomach (39%–68%), and is largely associated with *SMAD4* pathogenic variants.⁹⁴ Excess risk of nongastrointestinal cancer is not reported in JPS.

PTEN-HAMARTOMA TUMOR SYNDROME

Clinical Features

PHTS includes a variety of phenotypic variations known as CS, BRRS, and Proteus syndrome. The clinical diagnosis of the PHTS is made in patients meeting the revised diagnostic criteria, which include the presence of hamartomas of the skin and gastrointestinal tract (see Figures 1D and 2F–H), mucocutaneous lesions, macrocephaly, and an increased risk of benign and malignant lesions of the breast, thyroid, and endometrium.⁹⁵

Diagnosis

The genetic diagnosis of PHTS is established with a germline pathogenic variant in the phosphatase and tensin homolog (*PTEN*) gene. The clinical criteria for the diagnosis of CS is complex, and can be found at the National Comprehensive Cancer Network website.⁴⁹ The discussion for this article will focus on the manifestations and management of patients with a confirmed diagnosis of PHTS.

Genetics

The *PTEN* gene encodes a dual-function phosphatase that negatively regulates the growth-promoting activity of the phosphatidylinositol 3-kinase pathway. The PHTS family of syndromes are caused by inactivating pathogenic variants in *PTEN*, making it a classic tumor suppressor gene.⁹⁶ Germline pathogenic variants in the *PTEN* gene lead to heterogeneous phenotypes called PHTS, which includes CS, BRRS, and some cases of the *PTEN*-related Proteus syndrome (not considered further here).^{97,98} Germline pathogenic variants in *PTEN* are associated with other systemic nonpolyposis phenotypes, including abnormalities of the central nervous system and skeleton, but these will not be discussed here in detail. Patients with CS are also at risk for dysplastic cerebellar gangliocytoma (Lhermitte-Duclos disease), but the specific genetic basis of this and the other protean manifestations of

germline pathogenic variants in *PTEN* are not yet understood. Interestingly, gain-of-function germline pathogenic variants in the *WWP1* gene, an E3 ubiquitin ligase commonly up-regulated in cancers, inhibits the activity of *PTEN*, causes a CS-like syndrome (oligopolyposis and cancer-prone phenotype), and provides some insight into what may be responsible for PHTS in the absence of germline variants in *PTEN*.⁹⁹

Cancer Risk

Question: Which organs should undergo surveillance for cancer when caring for a patient with *PTEN* hamartoma tumor syndrome?

Recommendation: In *PTEN* hamartoma tumor syndrome, patients are at increased risk for cancer in multiple organs, including cancer of the breast, thyroid, kidney, uterus, colon, and skin.

Given this risk, we recommend a multidisciplinary approach to cancer surveillance in these organs. (Strong recommendation, low quality evidence)

Cancer risks for patients with germline pathogenic variants in *PTEN* are very high. The International Cowden Consortium reported in 2012 on 368 individuals with germline pathogenic variants in *PTEN*. Elevated standardized incidence ratios (SIRs) were found for carcinomas of the breast (SIR, 25.4; 95% CL, 19.8 and 32.0), thyroid (SIR, 51.1; 95% CL, 38.1 and 67.1), endometrium (SIR, 42.9; 95% CL, 28.1 and 62.8), colorectum (SIR, 10.3; 95% CL, 5.6 and 17.4), kidney (SIR, 30.6; 95% CL, 17.8 and 49.4), and melanoma (SIR, 8.5; 95% CL, 4.1 and 15.6). This led to cumulative lifetime risks for cancer of the breast at 85.2% (95% CL, 71.4% and 99.1%), thyroid 35.2% (95% CL, 19.7% and 50.7%), endometrium 28.2% (95% CL, 17.1% and 39.3%), colorectum 9.0% (95% CL, 3.8% and 14.1%), kidney 33.6% (95% CL, 10.4% and 56.9%), and melanoma 6% (95% CL, 1.6% and 9.4%).¹⁰⁰

A multicenter study from France estimated cancer risks in 154 patients with a germline pathogenic variant in *PTEN*.¹⁰¹ SIRs for female breast cancer were 39.1 (95% confidence interval [CI], 24.8–58.6), thyroid cancer 43.2 (95% CI, 19.7–82.1) in women and 199.5 (95% CI, 106.39–342.03) in men, melanoma 28.3 (96% CI, 7.6–35.4) in women and 39.4 (95% CI, 10.6–100.9) in men, and endometrial cancer 48.7 (95% CI, 9.8–142.3). Estimated cumulative lifetime risks by age 70 years were 85% for any cancer, 77% for female breast cancer, and 38% for thyroid cancer. Median age for developing cancer was 36 years. Although 85% were reported to have colonic polyps of any variety, risks for gastrointestinal cancer were not reported to be elevated in this series. However, a review of 211 patients with CS from the Mayo Clinic, including a review of the literature, reported a 16% lifetime risk for CRC. In this study, only 46% had a proven germline pathogenic variant in *PTEN*, and the cumulative risk for any cancer by age 70 years was 89%.¹⁰²

In a study of 2548 patients who met “relaxed” International Cowden’s Consortium criteria for CS with 5 or more gastrointestinal polyps in which at least 1 was considered hyperplastic or hamartomatous, germline pathogenic variants in *PTEN* were found in 127 (5%).¹⁰³ At endoscopy, gastrointestinal polyps were found throughout the gut, and one-half were considered “hyperplastic.” In this group, 13% of those who underwent colonoscopy developed CRC (7.1% of the entire series of patients), all before age 50 years (the youngest was 35 years old), with a SIR for CRC of 224.1 (95% CI, 109.3–411.3).

Question: Which gastrointestinal findings should prompt a genetic evaluation for *PTEN* hamartoma tumor syndrome?

Recommendation: We recommend individuals with multiple gastrointestinal hamartomas or ganglioneuromas undergo genetic evaluation for Cowden’s syndrome and related conditions. (Strong recommendation, low quality of evidence)

A multicenter study from 9 countries of 180 carriers of germline pathogenic variants in *PTEN* estimated the cumulative risk of any cancer or Lhermitte-Duclos disease by age 60 years was 56% for men and 87% for women.¹⁰⁴ Increased risk was reported for cancers of the breast, thyroid, endometrium, skin, kidneys, colorectum, and lungs. An earlier report from this group on 156 patients reported that benign gastrointestinal polyps were found in 31% of patients at a mean age of 38 years (range, 18–62 years) and most were considered “hamartomas.”¹⁰⁵ The cumulative risk of developing polyps was 70% by age 60 years. The cumulative risk of developing CRC was 18% by age 60 years (occurring between ages 53 and 62 years), suggesting that surveillance colonoscopy might not be necessary in early adult life.

Gastrointestinal Polyps and Polyposis

Gastrointestinal polyps are frequently found in patients with germline pathogenic variants in *PTEN*, with variable prevalence. Estimates suggest that 35%–93% of patients with CS have gastrointestinal polyps, but in some reports the germline basis of the disease was not known and a uniform interpretation of non-adenomatous polyps had not been established.^{103,106} The wide range of estimated polyp prevalence is probably a reflection of the heterogeneity of the disease and challenges in making definitive diagnoses outside of germline tests. Gastrointestinal polyps include hyperplastic polyps, inflammatory polyps, ganglioneuromas, lipomas, adenomas, and the nonspecific term hamartomas.

A prospective study of 127 *PTEN* pathogenic variant carriers from 2548 subjects in an International Cowden Consortium study in which 69 underwent endoscopic studies found that 64 (93%) had gastrointestinal polyps.¹⁰³ Of that group, one-half had hyperplastic polyps, but all of the above-listed polyps (plus adenomas) were also found. The number of polyps ranged from 1 to “innumerable,” and were distributed throughout the gut.

BRRS is caused by germline pathogenic variants in the *PTEN* gene.^{98,107} It is a pediatric condition associated with macrocephaly, developmental delay, gastrointestinal hamartomatous polyps, and pigmented macules on the toes and glans penis. It has also been called the Bannayan-Zonana or Ruvalcaba-Riley-Smith syndrome. Inexplicably, members of the same family may have features of either BRRS or CS,¹⁰⁸ and some cases of BRRS do not have detectable pathogenic variants in *PTEN*.¹⁰⁹ Confusing overlap between CS and JPS can occur when, as mentioned above, a chromosomal deletion occurs on chromosome 10q22.3–q24.1. This leads to the loss of both *PTEN* and *BMPRIA*²⁸; a situation that can also look like BRRS, also known as juvenile polyposis of infancy (see section on JPS).^{110,111}

In a smaller study of 19 patients with CS (in which only 12 were shown to have germline pathogenic variants in *PTEN*), pan-colonic polyps were found in 58%, pan-gastrointestinal polyps in 42%, and the pathological interpretation of the polyps included inflammatory polyps in 95%, but also adenomas, lipomas, and ganglioneuromas.¹¹² Moreover, there was more than 1 pathological variety in 79% of patients, indicating the clinical heterogeneity in this entity. Esophageal glycogenic acanthosis is also found in PHTS, which is a benign condition and there is no reported increased risk for esophageal cancer.¹¹³

Surveillance of Affected Individuals

Surveillance recommendations are provided in Table 4. The assessment at diagnosis of CS/PHTS should include a complete (and especially dermatologic and neurologic) clinical examination,

mammography and breast MRI, thyroid ultrasound, transvaginal ultrasound, upper gastrointestinal endoscopy, colonoscopy, and renal ultrasound. Although there are no data regarding risk-reduction surgery in women with CS, the option for risk-reducing mastectomy and hysterectomy should be discussed on a case-by-case basis.

The risk of CRC in patients with CS is estimated to be up to 9%–18% lifetime risk with mean age at diagnosis of 44 years, but ranging from 35 to 49 years.^{100,102,103,105,114,115} Consequently, expert opinion recommends colonoscopy starting at age 35 years unless symptomatic or if a close relative has had colon cancer before age 40 years, then start 10 years before the earliest known colon cancer in the family and repeat every 5 years or more frequently if the patient is symptomatic or polyps are found. This recommendation differs from a recent American College of Gastroenterology Guideline,³² which recommends initiating colonoscopy at age 15 years. Recent evidence suggesting later onset of significant colon cancer risk informed our recommendations.

Colectomy is rarely required in patients with CS and is reserved for patients with CRC or in whom surveillance and clearing of premalignant lesions is not endoscopically feasible.

Question: What is the recommended colonoscopic surveillance in individuals identified with *PTEN* hamartoma tumor syndrome?

Recommendation: We suggest colonoscopy surveillance to begin at age 35 years (or 10 years younger than age of any relative with colorectal cancer), repeated at intervals no greater than 5 years, depending on polyp burden. (Weak recommendation, low quality of evidence)

Breast Cancer

The management of breast cancer risk begins at age 25 years with clinical breast examination every 6–12 months; annual mammography and breast MRI surveillance starting between ages 30 and 35 years or 5–10 years before the earliest known breast cancer in the family.⁴⁹ In patients with PHTS, referral to a breast cancer specialist for management of breast cancer surveillance is reasonable and a multidisciplinary approach, including a breast surgeon, is recommended when prophylactic mastectomy is being considered.

Endometrial Cancer

The management of endometrial cancer risk begins between the ages of 30 and 35 years. Endometrial cancer can often be detected early on the basis of symptoms, and women should be educated regarding seeking medical attention on the basis of symptoms, including abnormal uterine bleeding or postmenopausal bleeding. Endometrial biopsy is sensitive and specific for endometrial cancer and surveillance via biopsy every 1–2 years can be considered.^{33,49}

Thyroid Cancer

An annual thyroid ultrasound should be performed, beginning at the time of PHTS diagnosis (including in childhood).^{33,49}

Renal Cancer

The recommended management of kidney cancer risk is an annual renal ultrasound and/or renal MRI starting at age 40 years if there is a family history of renal cancer, or every 2 years if not.^{33,49}

Table 5. Areas Requiring Further Investigation in Hamartomatous Polyposis Syndromes

1. Discover the germline variants in families with clinically recognizable syndromes who do not have germline mutations in the genes known to be associated with these syndromes.
2. Understand the effects of haplo-insufficiency of <i>STK11</i> in PJS and the <i>SMAD4</i> in JPS on the development of polyposis.
3. Identify safe and effective pharmacological interventions for the inhibition of polyp formation children and adults with the hamartomatous polyposis syndromes.
4. Determine whether pharmacological intervention can mitigate cancer risk in PJS and JPS in adults (independent of the polyposis risk).
5. Develop simulation models in adults with a hamartomatous polyposis syndrome to determine the optimal endoscopic frequency.
6. Design and implement studies to determine the best imaging modalities and surveillance intervals in adults with a hamartomatous polyposis syndrome.
7. Determine whether the hamartomatous phenotype in pediatric patients predicts the cancer risk in adulthood.
8. Perform outcomes studies evaluating the transition of care in pediatric polyposis patients entering adult life, including outcomes such as compliance with surveillance.
9. Determine difference in phenotype and cancer risk in individuals who meet clinical criteria, but do not have an identifiable pathogenic variant compared to those with a germline pathogenic variant in a hamartomatous polyposis syndrome.
10. Determine the lifetime cancer risks in patients with a pathogenic variant, but without clinical manifestations (those identified incidentally on multigene panel testing).
11. Quantify the frequency of mosaicism as a cause of the hamartomatous polyposis syndromes.
12. Determine the optimal modality of small bowel evaluation in individuals with PJS.

Melanoma

The management of melanoma risk includes an annual clinical skin examination beginning at age 18 years.³³

Summary

Germline pathogenic variants in *PTEN* are associated with variety of gastrointestinal hamartomatous polyps and a markedly increased risk of cancer, but largely in nongastrointestinal organs. There is a moderate increase in the risk of CRC, but it may be that the age of risk is such that colonoscopic surveillance can be withheld until age 35 years or later or 10 years younger than the age of the youngest relative with CRC. However, the reports of CRC risk have been variable,^{95,116} and no surveillance recommendation has been rigorously evaluated. The principal organs at risk for cancer include the breast, thyroid, endometrium, and colorectum.

HEREDITARY MIXED POLYPOSIS SYNDROME

The Genetic Basis of the Disease and Cancer Risk

HMPS is a rare autosomal dominant disease reported in only a few families. It is characterized by attenuated colonic polyposis. The polyps include adenomas, hyperplastic, and a particular

polyp with an admixture of variable histologies including adenomatous, hyperplastic, juvenile, and mixed polyps.¹¹⁷ In the original HMPS kindred described by Whitelaw et al,¹¹⁸ 13 members were diagnosed with CRC and 23 developed multiple polyps of several different histologic types (adenomatous, hyperplastic, and juvenile). HMPS has been associated with large duplications of the promoter region or entire *GREM1* gene.^{119–121} These unusual pathogenic variants increase the expression of the gene product, which then inhibits the bone morphogenetic protein (BMP) pathway.¹¹⁹ This promoter duplication was found in 0.7% of Ashkenazi Jews in Israel who met clinical criteria for Lynch syndrome.¹²⁰ The largest series reported 4 families with 16 affected members; the onset of polyposis starts in the late 20s, which is when colonoscopic surveillance should begin.¹²² There are not enough data to know the optimal surveillance intervals or whether extraintestinal neoplasia is a risk. The underlying genetic basis of the majority of HMPS families is unknown.

FUTURE RESEARCH CONSIDERATIONS

The hamartoma syndromes are rare and it is difficult for single institutions or even collaborative centers to accumulate enough cases and follow them prospectively long enough to develop robust conclusions about cancer risk. Modeling, simulation, and collaborative multicenter clinical studies can be used to help clarify the benefits and risks of various interventions and surveillance programs. The management of these diseases changes dramatically when the patient matures from childhood to adulthood—which is where almost all of the cancer risk lies. Important knowledge gaps are listed in Table 5 and expanded upon below.

Peutz-Jeghers Syndrome

As indicated above, *STK11/LKB1* pathogenic variants result in up-regulation of inflammatory cytokines and promotion of overgrowth of both stromal and normal gastrointestinal epithelium, driving polyp formation (Table 5). Up-regulation of cytokines is associated with hyperactivation of JAK-STAT, which contributes to inflammation and cancer. Inhibition of JAK-STAT significantly reduced polyp growth in mice. Consequently, the JAK inhibitor ruxolitinib (already clinically approved for myeloproliferative disease) may have therapeutic potential in patients with PJS.²¹ Also, LKB1 activity inhibits the activation of adenosine monophosphate-dependent protein kinase. Loss of LKB1 activity results in adenosine monophosphate-dependent protein kinase activation with up-regulation of mTORC1 signaling contributing to growth of PJS polyposis. Targeting of adenosine monophosphate-dependent protein kinase activation is currently being investigated.¹²³

Juvenile Polyposis Syndrome

Of note is the mouse model of juvenile polyposis created by a conditional knockout of the *SMAD4* gene only in T-cell populations.⁷³ These animals spontaneously developed massive polyps within the gastroduodenal region. The epithelium in the polyps contained increased expression of pro-inflammatory cytokines, particularly interleukin-11, a cytokine known to promote gastric epithelial cell survival and hyperplasia with evidence of increased TH17 cell activity. This suggests possible therapeutic targets for the future chemoprevention of juvenile polyposis.

SUMMARY AND CONCLUSIONS

Among the gastrointestinal hamartomatous polyposis syndromes, PJS is the best understood. The polyps are readily identified as PJ polyps pathologically, and only 1 known gene (*STK11*) is associated with this entity. The phenotype is clinically distinct. Not all patients (or families) have detectable germline pathogenic variants in *STK11*, so additional genes in this pathway are possibly contributors to this entity. The polyps may evolve through an expansion of elements of the gut stroma due to haploinsufficiency of *STK11*. In children, the main risks are for gastrointestinal obstruction and bleeding. Later in adult life, a very high risk of intestinal and extraintestinal cancers exists. The major organs at risk for cancer (in order of decreasing relative risk) include the small intestine, stomach, pancreas, colon, esophagus, ovary, lung, uterus, and breast, with estimated lifetime risks for any cancer reaching >90%. The ovaries and testes are also at risk for rare variant tumors. The screening and surveillance strategies change dramatically when the children reach adulthood.

JPS has principal risks for obstruction and bleeding in the pediatric ages, but the management in adults shifts to cancer risks in the stomach and colon. Importantly, unlike PJS, the gastrointestinal cancers tend to arise within the juvenile polyps, suggesting that polyp removal might prevent cancer. Cancer risk is linked to germline pathogenic variants in the *SMAD4* gene rather than the *BMPRIA* gene. The polyps are histologically inflammatory in nature, and their growth may be driven by haploinsufficiency of *SMAD4* in the immune cells of the polyp stroma. The management focuses on the polyps in children and cancers later in life, when endoscopic approaches are the mainstay of surveillance.

The hamartoma syndromes associated with germline pathogenic variants in *PTEN* raise a completely different clinical challenge. Although nearly all patients with PHTS have a variety of different hamartomatous gastrointestinal polyps, studies suggest an increased risk of cancer in this setting.^{100,103} The extracolonic cancer risks are greatest for the breast, thyroid, melanoma, and endometrium. The absolute risk for CRC appears to be increased, ranging from 9% to 18% during a lifetime, and the rest of the gastrointestinal tract does not have an established increase in cancer risk. The age for beginning surveillance for CRC remains to be determined, but the early onset of CRC provides some suggestions. Be aware that the *PTEN* gene and *BMPRIA* are located near one another on chromosome 10q, and that large-scale chromosomal deletions could adversely affect both genes, complicating the clinical picture. Long-term prospective studies of mutation carriers are still needed to further clarify the risk of cancer and the role of surveillance in these syndromes. With increases in genetic testing and evaluation, future studies will be conducted with more robust cohorts of genetically characterized, less heterogeneous populations. However, there is also a need to study patients and families with unusual phenotypes where no genotype can be found.

Conflicts of Interest

The authors disclose no conflicts of interest relative to the current work since 2016. These authors disclose the following industry relationships (consulting, research, reimbursement) without conflict of interest relevant to the current work since 2016: C. Richard Boland: Ambry Genetics. Gregory Idos: Myriad Genetics, Inc. Carol Burke: Salix Pharmaceuticals, Ferring Pharmaceuticals, Aries Pharmaceuticals, Pfizer, Cancer Prevention Pharmaceuticals, Janssen Pharmaceuticals, SLA Pharma AG, and Freenome Holdings, Inc. Samir Gupta: Freenome Holdings, Inc, Guardant Health, Inc, CellMax, Inc, and Mallinckrodt Pharmaceuticals. Brian C. Jacobson:

Motus, GI, Dark Canyon LLC, and Remedy Partners. Aasma Shaukat: Iterative Scopes, Freenome Holdings Inc. Swati G. Patel: Olympus America, Freenome Holdings Inc, ERBE USA. Sapna Syngal: Myriad Genetics, Inc, DC Health, Inc, and GlaxoSmith Kline, Inc. Douglas Robertson: Covidien, Freenome Holdings, Inc, and Amadix. The remaining authors disclose no conflicts.

REFERENCES

- Guyatt GH, Oxman AD, Kunz R, et al. What is "quality of evidence" and why is it important to clinicians? *BMJ* 2008;336:995–998.
- Lu KH, Wood ME, Daniels M, et al. American Society of Clinical Oncology Expert Statement: collection and use of a cancer family history for oncology providers. *J Clin Oncol* 2014;32:833–840.
- Hampel H, Bennett RL, Buchanan A, et al. A practice guideline from the American College of Medical Genetics and Genomics and the National Society of Genetic Counselors: referral indications for cancer predisposition assessment. *Genet Med* 2015;17:70–87.
- Heald B, Hampel H, Church J, et al. Collaborative Group of the Americas on Inherited Gastrointestinal Cancer Position statement on multigene panel testing for patients with colorectal cancer and/or polyposis. *Fam Cancer* 2020;19:223–239.
- Attard TM, Burke CA, Hyer W, et al. ACG clinical report and recommendations on transition of care in children and adolescents with hereditary polyposis syndromes. *Am J Gastroenterol* 2021;116:638–646.
- van Lier MG, Wagner A, Mathus-Vliegen EM, et al. High cancer risk in Peutz-Jeghers syndrome: a systematic review and surveillance recommendations. *Am J Gastroenterol* 2010;105:1258–1264; author reply 1265.
- Latchford AR, Phillips RK. Gastrointestinal polyps and cancer in Peutz-Jeghers syndrome: clinical aspects. *Fam Cancer* 2011;10:455–461.
- Shaco-Levy R, Jasperson KW, Martin K, et al. Morphologic characterization of hamartomatous gastrointestinal polyps in Cowden syndrome, Peutz-Jeghers syndrome, and juvenile polyposis syndrome. *Hum Pathol* 2016;49:39–48.
- Menko FH. LKB1/STK11, Peutz-Jeghers syndrome and cancer. *Fam Cancer* 2011;10:413–414.
- Jass JR. Colorectal polyposis: from phenotype to diagnosis. *Pathol Res Pract* 2008;204:431–447.
- Beggs AD, Latchford AR, Vasen HF, et al. Peutz-Jeghers syndrome: a systematic review and recommendations for management. *Gut* 2010;59:975–986.
- Amos CI, Bali D, Thiel TJ, et al. Fine mapping of a genetic locus for Peutz-Jeghers syndrome on chromosome 19p. *Cancer Res* 1997;57:3653–3656.
- Hemminki A, Markie D, Tomlinson I, et al. A serine/threonine kinase gene defective in Peutz-Jeghers syndrome. *Nature* 1998;391:184–187.
- Hardie DG. The LKB1-AMPK pathway—friend or foe in cancer? *Cancer Cell* 2013;23:131–132.
- Shaw RJ, Bardeesy N, Manning BD, et al. The LKB1 tumor suppressor negatively regulates mTOR signaling. *Cancer Cell* 2004;6:91–99.
- Knudson AG Jr. Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci USA* 1971;68:820–823.
- Hernan I, Roig I, Martin B, et al. De novo germline mutation in the serine-threonine kinase *STK11/LKB1* gene associated with Peutz-Jeghers syndrome. *Clin Genet* 2004;66:58–62.
- Katajisto P, Vaahtomeri K, Ekman N, et al. LKB1 signaling in mesenchymal cells required for suppression of gastrointestinal polyposis. *Nat Genet* 2008;40:455–459.
- Poffenberger MC, Metcalfe-Roach A, Aguilar E, et al. LKB1 deficiency in T cells promotes the development of gastrointestinal polyposis. *Science* 2018;361:406–411.
- Ollila S, Domenech-Moreno E, Laajanen K, et al. Stromal Lkb1 deficiency leads to gastrointestinal tumorigenesis involving the IL-11-JAK/STAT3 pathway. *J Clin Invest* 2018;128:402–414.
- Hollstein PE, Shaw RJ. Inflamed T cells and stroma drive gut tumors. *Science* 2018;361:332–333.
- Amos CI, Keitheri-Cheteri MB, Sabripour M, et al. Genotype-phenotype correlations in Peutz-Jeghers syndrome. *J Med Genet* 2004;41:327–333.
- Boron P, De Rosa M, Nedoszytko B, et al. Specific Alu elements involved in a significant percentage of copy number variations of the *STK11* gene in patients with Peutz-Jeghers syndrome. *Fam Cancer* 2015;14:455–461.

24. Zbuk KM, Eng C. Hamartomatous polyposis syndromes. *Nat Clin Pract Gastroenterol Hepatol* 2007;4:492–502.
25. Idrees MT, Ulbright TM, Oliva E, et al. The World Health Organization 2016 classification of testicular non-germ cell tumours: a review and update from the International Society of Urological Pathology Testis Consultation Panel. *Histopathology* 2017;70:513–521.
26. Giardiello FM, Brensinger JD, Tersmette AC, et al. Very high risk of cancer in familial Peutz-Jeghers syndrome. *Gastroenterology* 2000;119:1447–1453.
27. Resta N, Pierannunzio D, Lenato GM, et al. Cancer risk associated with STK11/LKB1 germline mutations in Peutz-Jeghers syndrome patients: results of an Italian multicenter study. *Dig Liver Dis* 2013;45:606–611.
28. Hearle N, Schumacher V, Menko FH, et al. Frequency and spectrum of cancers in the Peutz-Jeghers syndrome. *Clin Cancer Res* 2006;12:3209–3215.
29. van Lier MG, Westerman AM, Wagner A, et al. High cancer risk and increased mortality in patients with Peutz-Jeghers syndrome. *Gut* 2011;60:141–147.
30. Lim W, Olschwang S, Keller JJ, et al. Relative frequency and morphology of cancers in STK11 mutation carriers. *Gastroenterology* 2004;126:1788–1794.
31. Mehenni H, Resta N, Park JG, et al. Cancer risks in LKB1 germline mutation carriers. *Gut* 2006;55:984–990.
32. Chen HY, Jin XW, Li BR, et al. Cancer risk in patients with Peutz-Jeghers syndrome: a retrospective cohort study of 336 cases. *Tumour Biol* 2017;39(6):1010428317705131.
33. Syngal S, Brand RE, Church JM, et al. ACG clinical guideline: genetic testing and management of hereditary gastrointestinal cancer syndromes. *Am J Gastroenterol* 2015;110:223–262; quiz 263.
34. Gupta S, Provenzale D, Regenbogen SE, et al. NCCN Guidelines Insights: Genetic/Familial High-Risk Assessment: Colorectal, Version 3.2017. *J Natl Compr Cancer Netw* 2017;15:1465–1475.
35. Giardiello FM, Trimbath JD. Peutz-Jeghers syndrome and management recommendations. *Clin Gastroenterol Hepatol* 2006;4:408–415.
36. Utsunomiya J, Gocho H, Miyanaga T, et al. Peutz-Jeghers syndrome: its natural course and management. *Johns Hopkins Med J* 1975;136:71–82.
37. Hinds R, Philp C, Hyer W, et al. Complications of childhood Peutz-Jeghers syndrome: implications for pediatric screening. *J Pediatr Gastroenterol Nutr* 2004;39:219–220.
38. Fallon SC, Lopez ME, Zhang W, et al. Risk factors for surgery in pediatric intussusception in the era of pneumatic reduction. *J Pediatr Surg* 2013;48:1032–1036.
39. Gastineau S, Viala J, Caldari D, et al. Contribution of capsule endoscopy to Peutz-Jeghers syndrome management in children. *Dig Liver Dis* 2012;44:839–843.
40. Bizzarri B, Borrelli O, de'Angelis N, et al. Management of duodenal-jejunal polyps in children with peutz-jeghers syndrome with single-balloon enteroscopy. *J Pediatr Gastroenterol Nutr* 2014;59:49–53.
41. Latchford AR, Neale K, Phillips RK, et al. Peutz-Jeghers syndrome: intriguing suggestion of gastrointestinal cancer prevention from surveillance. *Dis Colon Rectum* 2011;54:1547–1551.
42. Tomlinson IP, Houlston RS. Peutz-Jeghers syndrome. *J Med Genet* 1997;34:1007–1011.
43. Goverde A, Korsse SE, Wagner A, et al. Small-bowel surveillance in patients with Peutz-Jeghers syndrome: comparing magnetic resonance enteroclysis and double balloon enteroscopy. *J Clin Gastroenterol* 2017;51:e27–e33.
44. McGrath DR, Spigelman AD. Preventive measures in Peutz-Jeghers syndrome. *Fam Cancer* 2001;1:121–125.
45. Ohmiya N, Taguchi A, Shirai K, et al. Endoscopic resection of Peutz-Jeghers polyps throughout the small intestine at double-balloon enteroscopy without laparotomy. *Gastrointest Endosc* 2005;61:140–147.
46. Oncel M, Remzi FH, Church JM, et al. Benefits of 'clean sweep' in Peutz-Jeghers patients. *Colorectal Dis* 2004;6:332–335.
47. van Lier MG, Mathus-Vliegen EM, Wagner A, et al. High cumulative risk of intussusception in patients with Peutz-Jeghers syndrome: time to update surveillance guidelines? *Am J Gastroenterol* 2011;106:940–945.
48. Hamdi Y, Soucy P, Kuchenbaeker KB, et al. Association of breast cancer risk in BRCA1 and BRCA2 mutation carriers with genetic variants showing differential allelic expression: identification of a modifier of breast cancer risk at locus 11q22.3. *Breast Cancer Res Treat* 2017;161:117–134.
49. Daly MB, Pilarski R, Yurgelun MB, et al. NCCN Guidelines Insights: Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic, Version 1.2020. *J Natl Compr Canc Netw* 2020;18:380–391.
50. Manahan ER, Kuerer HM, Sebastian M, et al. Consensus Guidelines on Genetic Testing for Hereditary Breast Cancer from the American Society of Breast Surgeons. *Ann Surg Oncol* 2019;26:3025–3031.
51. Korsse SE, Harinck F, van Lier MG, et al. Pancreatic cancer risk in Peutz-Jeghers syndrome patients: a large cohort study and implications for surveillance. *J Med Genet* 2013;50:59–64.
52. Chen KT. Female genital tract tumors in Peutz-Jeghers syndrome. *Human Pathol* 1986;17:858–861.
53. Canto MI, Harinck F, Hruban RH, et al. International Cancer of the Pancreas Screening (CAPS) Consortium summit on the management of patients with increased risk for familial pancreatic cancer. *Gut* 2013;62:339–347.
54. Canto MI, Almario JA, Schulick RD, et al. Risk of neoplastic progression in individuals at high risk for pancreatic cancer undergoing long-term surveillance. *Gastroenterology* 2018;155:740–751.e2.
55. Goggins M, Overbeek KA, Brand R, et al. Management of patients with increased risk for familial pancreatic cancer: updated recommendations from the International Cancer of the Pancreas Screening (CAPS) Consortium. *Gut* 2020;69:7–17.
56. McGowan L, Young RH, Scully RE. Peutz-Jeghers syndrome with "adenoma malignum" of the cervix. A report of two cases. *Gynecol Oncol* 1980;10:125–133.
57. Crispo A, Brennan P, Jockel KH, et al. The cumulative risk of lung cancer among current, ex- and never-smokers in European men. *Br J Cancer* 2004;91:1280–1286.
58. Pinsky PF, Zhu CS, Kramer BS. Lung cancer risk by years since quitting in 30+ pack year smokers. *J Med Screen* 2015;22:151–157.
59. Detterbeck FC, Mazzone PJ, Naidich DP, et al. Screening for lung cancer: diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143(Suppl):e78S–e92.
60. Wender R, Fontham ET, Barrera E Jr., et al. American Cancer Society lung cancer screening guidelines. *CA Cancer J Clin* 2013;63:107–117.
61. de Brabander J, Eskens F, Korsse SE, et al. Chemoprevention in patients with Peutz-Jeghers syndrome: lessons learned. *Oncologist* 2018;23:399.e33.
62. Udd L, Katajisto P, Rossi DJ, et al. Suppression of Peutz-Jeghers polyposis by inhibition of cyclooxygenase-2. *Gastroenterology* 2004;127:1030–1037.
63. Chow E, Macrae F. A review of juvenile polyposis syndrome. *J Gastroenterol Hepatol* 2005;20:1634–1640.
64. Schreiber IR, Baker M, Amos C, et al. The hamartomatous polyposis syndromes: a clinical and molecular review. *Am J Gastroenterol* 2005;100:476–490.
65. Latchford AR, Neale K, Phillips RK, et al. Juvenile polyposis syndrome: a study of genotype, phenotype, and long-term outcome. *Dis Colon Rectum* 2012;55:1038–1043.
66. Heald B, Rigelsky C, Moran R, et al. Prevalence of thoracic aortopathy in patients with juvenile polyposis syndrome-hereditary hemorrhagic telangiectasia due to SMAD4. *Am J Med Genet A* 2015;167A:1758–1762.
67. Aretz S, Stienen D, Uhlhaas S, et al. High proportion of large genomic deletions and a genotype phenotype update in 80 unrelated families with juvenile polyposis syndrome. *J Med Genet* 2007;44:702–709.
68. Aytac E, Sulu B, Heald B, et al. Genotype-defined cancer risk in juvenile polyposis syndrome. *Br J Surg* 2015;102:114–118.
69. Sweet K, Willis J, Zhou XP, et al. Molecular classification of patients with unexplained hamartomatous and hyperplastic polyposis. *JAMA* 2005;294:2465–2473.
70. Ngeow J, Heald B, Rybicki LA, et al. Prevalence of germline PTEN, BMP1A, SMAD4, STK11, and ENG mutations in patients with moderate-load colorectal polyps. *Gastroenterology* 2013;144:1402–1409, 1409.e1–e5.
71. Woodford-Richens K, Bevan S, Churchman M, et al. Analysis of genetic and phenotypic heterogeneity in juvenile polyposis. *Gut* 2000;46:656–660.
72. Alberici P, Gaspar C, Franken P, et al. Smad4 haploinsufficiency: a matter of dosage. *Pathogenetics* 2008;1(1):2.
73. Hahn JN, Falck VG, Jirik FR. Smad4 deficiency in T cells leads to the Th17-associated development of preterminal gastroduodenal lesions in mice. *J Clin Invest* 2011;121:4030–4042.

74. Kim BG, Li C, Qiao W, et al. Smad4 signalling in T cells is required for suppression of gastrointestinal cancer. *Nature* 2006;441:1015–1019.
75. Woodford-Richens K, Williamson J, Bevan S, et al. Allelic loss at SMAD4 in polyps from juvenile polyposis patients and use of fluorescence in situ hybridization to demonstrate clonal origin of the epithelium. *Cancer Res* 2000;60:2477–2482.
76. Blatter RH, Plasilova M, Wenzel F, et al. Somatic alterations in juvenile polyps from BMPR1A and SMAD4 mutation carriers. *Genes Chromosomes Cancer* 2015;54:575–582.
77. O'Malley M, LaGuardia L, Kalady MF, et al. The prevalence of hereditary hemorrhagic telangiectasia in juvenile polyposis syndrome. *Dis Colon Rectum* 2012;55:886–892.
78. Gallione CJ, Richards JA, Letteboer TG, et al. SMAD4 mutations found in unselected HHT patients. *J Med Genet* 2006;43:793–797.
79. Faughnan ME, Palda VA, Garcia-Tsao G, et al. International guidelines for the diagnosis and management of hereditary haemorrhagic telangiectasia. *J Med Genet* 2011;48:73–87.
80. Oliveira PH, Cunha C, Almeida S, et al. Juvenile polyposis of infancy in a child with deletion of BMPR1A and PTEN genes: surgical approach. *J Pediatr Surg* 2013;48:e33–e37.
81. Tsuchiya KD, Wiesner G, Cassidy SB, et al. Deletion 10q23.2-q23.33 in a patient with gastrointestinal juvenile polyposis and other features of a Cowden-like syndrome. *Genes Chromosomes Cancer* 1998;21:113–118.
82. Delnatte C, Sanlaville D, Mougenot JF, et al. Contiguous gene deletion within chromosome arm 10q is associated with juvenile polyposis of infancy, reflecting cooperation between the BMPR1A and PTEN tumor-suppressor genes. *Am J Hum Genet* 2006;78:1066–1074.
83. Menko FH, Kneepkens CM, de Leeuw N, et al. Variable phenotypes associated with 10q23 microdeletions involving the PTEN and BMPR1A genes. *Clin Genet* 2008;74:145–154.
84. Fox VL, Perros S, Jiang H, et al. Juvenile polyps: recurrence in patients with multiple and solitary polyps. *Clin Gastroenterol Hepatol* 2010;8:795–799.
85. Wu TT, Rezaei B, Rashid A, et al. Genetic alterations and epithelial dysplasia in juvenile polyposis syndrome and sporadic juvenile polyps. *Am J Pathol* 1997;150:939–947.
86. Brosens LA, van Hattem A, Hyllind LM, et al. Risk of colorectal cancer in juvenile polyposis. *Gut* 2007;56:965–967.
87. Gonzalez RS, Adsay V, Graham RP, et al. Massive gastric juvenile-type polyposis: a clinicopathological analysis of 22 cases. *Histopathology* 2017;70:918–928.
88. Friedl W, Uhlhaas S, Schulmann K, et al. Juvenile polyposis: massive gastric polyposis is more common in MADH4 mutation carriers than in BMPR1A mutation carriers. *Hum Genet* 2002;111:108–111.
89. Howe JR, Mitros FA, Summers RW. The risk of gastrointestinal carcinoma in familial juvenile polyposis. *Ann Surg Oncol* 1998;5:751–756.
90. Wirtzfeld DA, Petrelli NJ, Rodriguez-Bigas MA. Hamartomatous polyposis syndromes: molecular genetics, neoplastic risk, and surveillance recommendations. *Ann Surg Oncol* 2001;8:319–327.
91. MacFarland SP, Ebrahimzadeh JE, Zellek K, et al. Phenotypic differences in juvenile polyposis syndrome with or without a disease-causing SMAD4/BMPR1A variant. *Cancer Prev Res (Phila)* 2021;14:215–222.
92. Cohen S. Management of JPS in children and adolescents: a position paper from the ESPGHAN Polyposis Working Group. *J Pediatr Gastroenterol Nutr* 2019;68:453–462.
93. Faughnan ME, Mager JJ, Hetts SW, et al. Second international guidelines for the diagnosis and management of hereditary hemorrhagic telangiectasia. *Ann Intern Med* 2020;173:989–1001.
94. Campos FG, Figueiredo MN, Martinez CA. Colorectal cancer risk in hamartomatous polyposis syndromes. *World J Gastrointest Surg* 2015;7:25–32.
95. Pilarski R, Burt R, Kohlman W, et al. Cowden syndrome and the PTEN hamartoma tumor syndrome: systematic review and revised diagnostic criteria. *J Natl Cancer Inst* 2013;105:1607–1616.
96. Chalhoub N, Baker SJ. PTEN and the PI3-kinase pathway in cancer. *Annu Rev Pathol* 2009;4:127–150.
97. Liaw D, Marsh DJ, Li J, et al. Germline mutations of the PTEN gene in Cowden disease, an inherited breast and thyroid cancer syndrome. *Nat Genet* 1997;16:64–67.
98. Marsh DJ, Coulon V, Lunetta KL, et al. Mutation spectrum and genotype-phenotype analyses in Cowden disease and Bannayan-Zonana syndrome, two hamartoma syndromes with germline PTEN mutation. *Hum Mol Genet* 1998;7:507–515.
99. Lee YR, Yehia L, Kishikawa T, et al. WWP1 Gain-of-function inactivation of PTEN in cancer predisposition. *N Engl J Med* 2020;382:2103–2116.
100. Tan MH, Mester JL, Ngeow J, et al. Lifetime cancer risks in individuals with germline PTEN mutations. *Clin Cancer Res* 2012;18:400–407.
101. Bubien V, Bonnet F, Brouste V, et al. High cumulative risks of cancer in patients with PTEN hamartoma tumour syndrome. *J Med Genet* 2013;50:255–263.
102. Riegert-Johnson DL, Gleeson FC, Roberts M, et al. Cancer and Lhermitte-Duclos disease are common in Cowden syndrome patients. *Heredit Cancer Clin Pract* 2010;8(1):6.
103. Heald B, Mester J, Rybicki L, et al. Frequent gastrointestinal polyps and colorectal adenocarcinomas in a prospective series of PTEN mutation carriers. *Gastroenterology* 2010;139:1927–1933.
104. Nieuwenhuis MH, Kets CM, Murphy-Ryan M, et al. Cancer risk and genotype-phenotype correlations in PTEN hamartoma tumor syndrome. *Fam Cancer* 2014;13:57–63.
105. Nieuwenhuis MH, Kets CM, Murphy-Ryan M, et al. Is colorectal surveillance indicated in patients with PTEN mutations? *Colorectal Dis* 2012;14:e562–e566.
106. Stanich PP, Owens VL, Sweetser S, et al. Colonic polyposis and neoplasia in Cowden syndrome. *Mayo Clin Proc* 2011;86:489–492.
107. Arch EM, Goodman BK, Van Wesep RA, et al. Deletion of PTEN in a patient with Bannayan-Riley-Ruvalcaba syndrome suggests allelism with Cowden disease. *Am J Med Genet* 1997;71:489–493.
108. Hendriks YM, Verhallen JT, van der Smagt JJ, et al. Bannayan-Riley-Ruvalcaba syndrome: further delineation of the phenotype and management of PTEN mutation-positive cases. *Fam Cancer* 2003;2:79–85.
109. Carethers JM, Furnari FB, Zigman AF, et al. Absence of PTEN/MMAC1 germ-line mutations in sporadic Bannayan-Riley-Ruvalcaba syndrome. *Cancer Res* 1998;58:2724–2726.
110. Alimi A, Weeth-Feinstein LA, Stettner A, et al. Overlap of juvenile polyposis syndrome and Cowden syndrome due to de novo chromosome 10 deletion involving BMPR1A and PTEN: implications for treatment and surveillance. *Am J Med Genet A* 2015;167:1305–1308.
111. Hiljadnikova Bajro M, Sukarova-Angelovska E, Adelaide J, et al. A new case with 10q23 interstitial deletion encompassing both PTEN and BMPR1A narrows the genetic region deleted in juvenile polyposis syndrome. *J Appl Genet* 2013;54:43–47.
112. Shaco-Levy R, Jasperson KW, Martin K, et al. Gastrointestinal polyposis in cowden syndrome. *J Clin Gastroenterol* 2017;51:e60–e67.
113. Borowsky J, Setia N, Rosty C, et al. Spectrum of gastrointestinal tract pathology in a multicenter cohort of 43 Cowden syndrome patients. *Mod Pathol* 2019;32:1814–1822.
114. Ni Y, He X, Chen J, et al. Germline SDHx variants modify breast and thyroid cancer risks in Cowden and Cowden-like syndrome via FAD/NAD-dependant destabilization of p53. *Hum Mol Genet* 2012;21:300–310.
115. Kato M, Mizuki A, Hayashi T, et al. Cowden's disease diagnosed through mucocutaneous lesions and gastrointestinal polyposis with recurrent hematochezia, unrevealed by initial diagnosis. *Intern Med* 2000;39:559–563.
116. Stanich PP, Pilarski R, Rock J, et al. Colonic manifestations of PTEN hamartoma tumor syndrome: case series and systematic review. *World J Gastroenterol* 2014;20:1833–1838.
117. Plesec T, Brown K, Allen C, et al. Clinicopathological features of a kindred with SCG5-GREM1-associated hereditary mixed polyposis syndrome. *Hum Pathol* 2017;60:75–81.
118. Whitelaw SC, Murday VA, Tomlinson IP, et al. Clinical and molecular features of the hereditary mixed polyposis syndrome. *Gastroenterology* 1997;112:327–334.
119. Jaeger E, Leedham S, Lewis A, et al. Hereditary mixed polyposis syndrome is caused by a 40-kb upstream duplication that leads to increased and ectopic expression of the BMP antagonist GREM1. *Nat Genet* 2012;44:699–703.
120. Laitman Y, Jaeger E, Katz L, et al. GREM1 germline mutation screening in Ashkenazi Jewish patients with familial colorectal cancer. *Genet Res* 2015;97:e11.
121. Lewis A, Freeman-Mills L, de la Calle-Mustienes E, et al. A polymorphic enhancer near GREM1 influences bowel cancer risk through differential CDX2 and TCF7L2 binding. *Cell Rep* 2014;8:983–990.
122. Lieberman S, Walsh T, Schechter M, et al. Features of patients with hereditary mixed polyposis syndrome caused by duplication of GREM1 and implications for screening and surveillance. *Gastroenterology* 2017;152:1876–1880.e1.
123. Shackelford DB, Shaw RJ. The LKB1-AMPK pathway: metabolism and growth control in tumour suppression. *Nat Rev Cancer* 2009;9:563–575.